

GENUS EREUNETES.

99. **Semipalmated Sandpiper.** (Stint.)
(*Ereunetes pusillus*.)

Very abundant from about the twentieth of May to the end of the first week in June, and from the middle of July until the end of the first week in September.

The bulk of them go further north to breed, but I have always found some pairs here during the summer.

GENUS CALIDRIS.

100. **Sanderling.**
(*Calidris arenaria*)

Very abundant in spring and autumn along the lake shores.

Arrive about the tenth of May, depart in October.

Breeds in the far north.

GENUS LIMOSA.

101. **Marbled Godwit.**
(*Limosa fedoa*.)

A very rare visitor.

102. **Hudsonian Godwit.**
(*Limosa hæmastica*.)

Rather more frequently seen than the last, but still very rare.

GENUS TOTANUS.

SUBGENUS GLOTTIS.

103. **Greater Yellow-legs.**
(*Totanus melanoleucus*.)

Common in spring from early in April until the first week in May.

In the autumn sometimes abundant, arriving towards the end of July and remaining until the end of October.

Breeds in the far north.

104. **Yellow legs.**
(*Totanus flavipes*.)

Not common in spring, abundant in autumn, its movements in the spring correspond with those of the last species.

In the fall it arrives early in July and leaves before the middle of September.

Breeds in the far north.