



Fig. 3.



Fig. 4.

Fig. 3. The contents of a hernial sac in a pseudohermaphrodite. This mass was 7 inches long and 3 inches broad. It contained a considerable amount of turbid yellow fluid, a loop of small distended bowel, and the right testicle and epididymis. After the hernial ring had been cut, the bowel was reduced, but the testicle and epididymis were so firmly adherent to the sac, as noted in the picture, that it was necessary to remove them.

Fig. 4. Distribution of hair in a pseudohermaphrodite. The healed scar is readily seen. The pubic hair was well developed and extended upward a short distance above the umbilicus. The photograph does not show this very clearly. The legs were thickly covered with hair conforming closely to that of a male.

Leg (great troch.)	90 cm.	98 cm.	94 cm.	Nipples (center to center)	23.5 cm.	26 cm.	21 cm.
Circumference of chest	88.5 cm.	96 cm.	88 cm.	Nipple to umbilicus	28 cm.		
On inspiration			88 cm.	Sym. to spinous process of 5th lumbar vertebra			23.5 cm.
On expiration			84 cm.	Arm			
Width of shoulders	38.5 cm.	42.5 cm.		Length of humerus (axis of joints)			29 cm.
Center of			33 cm.	Length of forearm (axis of joints)			29 cm.
Humerus			32.5 cm.	Length of hand			27 cm.
Acromion process			49.5 cm.	Circumf. of humerus			27 cm.
Skin measurements				Circumf. of forearm			25 cm.
Width of waist	21 cm.	24 cm.	27 cm.	Leg			
Circum. of waist			74 cm.	Length of thigh (axis of joint)			44 cm.
Width of hips at crest of ilium	34.5 cm.	38 cm.	31 cm.	Length of thigh from ant. sup. sp. to bottom of knee-cap			51 cm.
Width at level of great trochanter	34 cm.	38 cm.	32 cm.	Length of leg			44 cm.
Pelvis between anterior superior spines	26.5 cm.	29.5 cm.	27 cm.				
Between post. sup. spines	12 cm.	13.5 cm.	11 cm. ¹				

¹ Angle with central skin fold 90° (sacral triangle).