

# NOTES ON SPECIALTIES.

## ANTISEPSIS OF THE INTESTINAL CANAL.

The griping pain and flatulence which accompany bowel and stomach complaints, particularly during the heated term, are so readily overcome and controlled by the timely administration of one or two Antikamnia & Salol Tablets, repeated every two or three hours, that it behooves us to call our readers' attention to the grand efficacy of this well-known remedy in these conditions. The above doses, are of course, those for adults. Children should be given one-fourth tablet for each five years of their age. When the attack is very severe, or when the disturbance is evidenced at or near the time of the menstrual period, we find it preferable to give two Antikamnia & Codeine Tablets, alternately with the Antikamnia & Salol Tablets. The latter Tablets promptly arrest excessive fermentation and have a pronounced sedative effect on the mucous membranes of the bowels and stomach, and will check the various diarrhœas without any untoward effect.



## WHEELER'S TISSUE PHOSPHATES.

The so-called peptonates and albuminates have to undergo in the organism before absorption complex chemical changes, in contradistinction to the inorganic mineral salts which are immediately absorbed as such, at a great saving of time and physiological energy. When a tonic and hæmatinic is indicated, therefore, it is best to prescribe Wheeler's Tissue Phosphates, a standard remedy with over a half-century of success behind it. A sample with literature will be sent free to any practitioner addressing

the T. B. Wheeler, M. D. Company,  
Montreal Canada.



## SAFE ANTISEPTICS IN GONORRHŒA

Tincture of iodine irrigations in solution of from one to four drachms to a quart of hot water is said to be one of the safest and best antiseptics that can be used in gonorrhœa. The strength of the solution and number of irrigations a day depend upon the stage of the disease. To keep the urine bland and non-irritating san-metto should be administered in teaspoonful doses three or four times daily throughout the treatment. In case of extreme acidity of the urine one of the potassium salts will be helpful.



## INFLAMMATORY DIARRHŒAS.

*By William Edward Fitch, M. D.,*

Lecturer on Surgery, Fordham University School  
of Medicine, New York City.

In discussing the subject we will speak of inflammation of the small and large intestines as a single disease. And without taking up the reader's valuable time in discussing etiology or symptomatology we will proceed at once to consider the medical treatment. The first step in this direction is to thoroughly evacuate the intestinal contents, and for this purpose no drug or combination of remedies has in our hands given the satisfaction that calomel has. Usually for a child of two years three grains are ordered rubbed up with sugar of milk and made into three powders and one administered every hour until all are taken, after which an old-fashioned dose of castor oil is given, which will produce several copious actions from