

improvement, and inside of a week his face was clear again. It is one of the greatest remedies I know of for those nasty, inflamed, raw looking and rapidly growing sores so often seen on school children's faces, and will cure them every time. I have prescribed it extensively in my practice for some years past, and always with success.—John Husson, M.D., 418 W. 124th Street, New York City.

THERE is no more important department of medical affairs than that of medical transfer. When a physician desires to sell his property and practice, it is of the utmost importance that it should be done with a minimum of publicity and a maximum of speed. The system adopted by Dr. Hamill, who conducts the Canadian Medical Exchange, is at once efficient and prompt, and offers every possible security to vendors, and we advise our readers to take advantage of his many years' experience when they are thinking of selling their practices. A partial list of the practices he has for sale will be found among our advertising columns every month, the complexion of which, of course, changes from time to time.

BATTLE & Co. have just issued the ninth of the series of twelve illustrations, of the Intestinal Parasites, and will send them free to physicians on application.

PROPER MEDICATION AND CHEERFUL COMPANY.—During the past two months, we have met with more la grippe than anything else, and the number of cases in which the pulmonary and bronchial organs have been very slightly or not at all involved, has been greater than we have noted in former invasions. On the contrary, grippal neuralgia, rheumatism and hepatitis have been of far greater frequency, while the nervous system has also been most seriously depressed. With each succeeding visitation of this trouble we have found it more and more necessary to watch out for the disease in disguise, and to treat these abnormal manifestations; consequently we have relied upon mild nerve sedatives, anodynes and tonics, rather than upon any specific line of treatment. Most cases will improve by being made to rest in bed and encouraging skin and kidney action, with possibly minute doses of blue pill or calomel. We have found much benefit from the use of antikamnia and salol tablets, two every three hours in the stage of pyrexia and muscular painfulness, and later on, when there was fever and bronchial cough and expectoration, from an antikamnia and codeine tablet every three