

Vertex slightly obtusely angled, the apex pointed, conical, half longer on middle than against the eyes, as long as the pronotum, slightly longer than its basal width, disc slightly sloping, anterior margin rounding to the full front except at the apex. Front and clypeus as in *belli*. Elytra long, inclined to be flaring posteriorly. Venation regular, but often obscured by numerous irregular reticulate veinlets on the clavus and in the ante-apical cells.

Colour.—Vertex pale, heavily washed with rusty brown, usually omitting a pale band before the eyes and often intensified as an oblique brown-dash either side the pointed apex. Face pale, with short brownish fuscous arcs and still darker sutures. Pronotum rusty or chocolate brown, with an anterior arcuated submarginal line paler. Elytra rusty or chocolate brown, with the nervures lighter, the irregular reticulations are usually strongly milky white and there is usually an oblique subhyaline light area beyond the middle of the costa. The apical cells smoky, with the nervures light.

Genitalia.—Female segment long, truncate or slightly concave posteriorly, with a broad angular median emargination, from which arises a strap-shaped tooth considerably exceeding the segment. Male valve triangular, apex round, plates strongly transversely convex, long, attenuate with the margins thickly beset with long hairs.

Described from four females and one male from San Francisco, collected by the author in September. The reticulate venation and white nervures will readily separate this species from any other described.

*Thamnotettix rupinata*, n. sp.

Form and general appearance of *intricata*, but lacking the supernumerary veinlets. Reddish or greenish brown, with a bisected black spot on the apex of vertex. Length, 5 mm.

Vertex slightly obtusely angular, the apex rounding, half longer on middle than against either eye, as wide as its median length in the female, narrower in the male, face long and narrow as in *intricata*. Elytra long and narrow, inclined to be closely appressed in the male, with the tips flaring, venation typical, regular, without extra veinlets.

Colour.—Vertex orange yellow, a large semi-circular black spot on the apex, bisected by the narrow, white median line, face pale, with narrow sutures and traces of arcs brown. Pronotum greenish or reddish brown, with a pale anterior margin, accentuated in the centre. Scutellum pale yellow, the basal angles olive. Elytra olive subhyaline with a reddish