

Dr. H. B. Anderson, Dr. W. P. Caven, Dr. E. E. King, Dr. John Amyot, Dr. F. N. G. Starr, Dr. R. D. Rudolf, Dr. R. J. Dwy. .

Section of Medicine.—Chairman, Dr. W. J. Wilson; Editor, Dr. John Ferguson; Secretary, Dr. W. Harley Smith. .

Section of Surgery.—Chairman, Dr. N. A. Powell; Editor, Dr. Shuttleworth; Secretary, Dr. H. A. Beatty.

Section of Pathology.—Chairman, Dr. W. Goldie; Editor, Dr. H. G. Hutchison; Secretary, Dr. Stanley Ryerson.

MEDICAL PREPARATIONS, ETC.

THE BEST HYPNOTIC.

A patient who would sleep but cannot sleep should be made to sleep. In the choice of a hypnotic the physician should always seek that one which not alone is most effective, but which presents the fewest disadvantages in the way of after effects. For years Bromidia has been the standard hypnotic prepared at the command of the profession. Through all the time that it has been known it has never failed in composition or efficiency. Its constituents have been of the purest, and in fact Bromidia has been the standard by which similar preparations have been measured. That the medical profession have appreciated its worth and thorough reliability is well apparent, from the place it holds in the regard of every physician who appreciates stability and honesty.—*The International Journal of Surgery*.

AN ANNUAL VISITOR.

We have just passed through our annual epidemic of la grippe, which, as usual, claimed its victims among all classes and conditions, mainly, however, among the classes where the resisting power was below par, or among sufferers from some chronic ailment. While the sequelæ and complications of this disease may assume almost any phase of acute inflammatory character, its primary effect is upon the nervous system. Therefore, we have no hesitancy in saying, no matter what the local inflammation may require as a medicine, by all means give antikamnia tablets as a nerve sedative and to relieve the muscular pains always present. We have seen a violent cough of bronchitis treated upon the general plan, with the cough as distressing at the end of twenty-four hours as at the beginning (promptly yield to six antikamnia tablets during an interval of six hours. La Grippe usually requires a double treatment, one directed to the influenza, and the other devoted to the complications present, be