

Of this number, cases 4, 5, 11, 12, 15, 16, 18, 20, 24, 25, 35, 48, 50, 52, 53, 54, 55, 57, 58, 72, 74, 79 were of more than ordinary interest, either clinically or on account of some unusual condition or complication.

In the 88 cases tuberculosis was found 30 times. These were classified as follows :—

CAUSE OF DEATH .....	11 times.
COMPLICATING .....	twice.
Latent Tuberculosis.....	7 times.
Obsolescent Tuberculosis.....	10 "

The clinical types in the fatal cases were :—

CHRONIC ULCERATIVE PULMONARY TUBERCULOSIS....	5 cases.
ACUTE TUBERCULOUS MENINGITIS.....	3 "
CHRONIC TUBERCULOUS PERITONITIS.....	1 case.
TUBERCULOUS TUMOUR OF BRAIN.....	1 "
GENITO-URINARY TUBERCULOSIS.....	1 "

Generalised arteriosclerosis was present in 11.36% of cases.

Cholelithiasis was met with in 5 cases, all in females.

The following ANOMALIES or MALFORMATIONS were noted :—

*Heart and Blood Vessels—*

Patent Foramen Ovale.....	12 times.
Hypoplasia of Aorta.....	once.

*Lungs—*

Abnormal lobation of right lung.....	10 times.
Abnormal lobation of left lung.....	3 "
Abnormal lobation of both lungs.....	once.

*Kidneys—*

Hypoplasia of left kidney.....	once.
Hypoplasia of both kidneys.....	"
Incomplete left double ureter.....	"
Complete right double ureter.....	"
Horseshoe kidney.....	twice.

*Liver—*

Liebermeister's grooves.....	6 times.
------------------------------	----------

*Spleen—*

Accessory spleens.....	16 times.
------------------------	-----------