

2nd Smith, Fairfield Plain. BANTAMS—golden Sebrights, 1st and 2nd McNeil, silver 1st and 2nd McNeil. Black breasted red, 1st Bonnick & Co. Toronto, 2nd Oldrieve Kingston. Golden Duckwing, 1st Bonnick & Co. 2nd Oldrieve. Silver 2d, Luxton, Red pyle, 1st Smith, 2nd Oldrieve. Brown red, 1st & 2nd Bonnick & Co. Japanese, 1st McNeil, 2nd Oldrieve, Pekin, 1st and 2nd McNeil. Any other variety, 1st and 2nd McNeil. TURKEYS—Bronze 1st and 2nd Main. White, 1st Smith. Any other variety, 1st Smith. GEESE—Bremen, 1st and 2nd Main. Toulouse, 1st Bogue 2nd McCormack, Rockton. Any other variety, 1st and 2nd Smith. DUCKS—Aylesbury, 1st and 2nd Bogue Rouen, 1st and 2nd Main. Pekin 1st Bogue, 2nd Cockburn jr. city. Cayuga, 1st and 2nd Luxton. GUINEA FOWL—1st Luxton, 2nd Smith, PEA FOWL—1st Luxton. MINORCAS—White 2nd Anderson, black 1st and 2nd McCormack. JAVAS—Black, 1st Luxton. ANDALUSIANS, 2nd Luxton.

CHICKS AND DUCKS 1889.

DORKINGS,—Colored 1st and 2nd Main. Silver-grey, 1st Bogue, 2nd Main. White, 1st and 2nd Bogue. COCHINS,—Buff, 1st McNeil, 2nd Bogue. Black, 1st McNeil, 2nd Bogue. Partridge, 1st Cockburn, city, 2nd Bogue. BRAHMAS,—Dark, 1st Peene, 2nd Luxton. Light, 1st Hooper, city, 2nd Clark & Climo. Hamburgs, Golden-spangled, 1st Bogue, 2nd McNeil. Silver, 1st and 2nd McNeil. Golden-pencilled, 1st McNeil, 2nd Oakhurst Jersey Farm. Silver, 1st Bogue, 2nd McNeil. Black, 1st McNeil, 2nd Luxton. LEGHORNS,—White, 1st Marshall, 2nd Edwards. Brown, 1st Dewar & Mitchell, 2d Stevens, Dunnville. Black, 1st and 2nd Hamilton. HOUDANS,—1st and 2nd Luxton. SPANISH,—Black, 1st Milton, 2nd Nicholson, city. PLYMOUTH ROCKS,—Barred, 1st and 2nd Clark & Climo. White, 1st and 2nd Clark & Climo. GAME,—Black-breasted red, 1st and 2nd Main. Silver Duckwing, 1st and 2nd Barber & Co. Red pyle, 1st Barber & Co. Brown breasted red, 1st and 2nd Barber & Co. Malays, 1st and 2nd Wardell. Pit, 1st Leggat. LANGSHANS,—1st McNeil, 2nd Alwin. MINORCAS,—Black, 1st McCormick, 2nd Marshall. POLANDS,—White crested, 1st McNeil, 2nd Bogue. Silver 1st Bogue. 2nd McNeil. Golden, 1st McNeil, 2nd Bogue. White bearded, 1st McNeil. JAVAS—Black, 1st and 2nd Luxton. BANTAMS—Black red game, 1st Barber & Co. 2nd Oldrieve, Kingston. Brown red 1st and 2nd Bonnick & Co. Silver duckwing, 1st Smith. Golden duckwing, 1st Aldrieve, Pyle, 1st Bonnick & Co., 2nd Barber & Co. SEBRIGHT—Golden, 1st and 2nd McNeil. Silver, 1st and 2nd McNeil. Japanese, 1st and 2nd McNeil. Pekin, 1st and 2nd McNeil.

Any other variety 1st and 2nd McNeil. DUCKLINGS—Aylesbury, 1st and 2nd Bogue. Rouen, 1st and 2nd Main. Pekin, 1st Bogue, 2nd Smith. Cayuga, 1st Luxton, 2nd Smith.

ANSWERS TO CORRESPONDENTS ON DISEASES, BREEDING, MANAGEMENT, ETC.

Correspondents are requested to make full use of this column. The answers to enquiries as to diseases will be answered by a well-known medical man and breeder. Please read the following rules carefully.

1. Give a concise, clear and exact statement of case, always stating age, sex, and breed.
2. Enclose 3 cents stamp for reply.
3. Report result, not necessarily for publication. *This is absolute.*
4. Acute cases requiring immediate treatment to be answered by mail in the first instance, later through POULTRY REVIEW for the benefit of our readers.
5. Write legibly and on one side of the paper only.
6. Answers to be to name in full, initials or *nom de plume*, the first preferred.

QUE.—I have a Game rooster, pure bred, two years and six months old, he has a swelling on the bottom of his right foot and it seems to be very sore. Another thing, when he stoops down to pick his food up off the ground saliva runs out of his mouth, please tell me what to do for him, and oblige.

Toronto.

F. H. FLINT.

ANS.—Your game cock has the disease known as "Bumble Foot," an inflammation that usually ends in the formation of an abscess (matter or pus). (1) Put the bird in a quiet place by himself. Feed him on soft food slightly peppered and only half the usual quantity for a couple of days. Give him a compound rhubarb pill for three days in succession. Give him also some green food chopped up fine. (2) Poultice the foot constantly until the swelling either goes down or bursts. Keep poulticing in the latter case till the foot gets quite soft and you can pick away any cheesy-like matter that is to be seen. When this is all away the foot will heal, when it is healed if it is still too large paint it every other day for two weeks, with Tincture of Iodine.

QUE.—I have a choice Buff Cochin Cock 1 year old. When he stands he can't stand long but always breaks down, it seems to me as if

he is weak in the legs, he has been that way about three weeks already, otherwise he is healthy, could you tell me anything to cure this. I would be very glad if you would tell me of something that would cure him.

Yours truly,

J. W. HUEHNERGARD.

Heidelberg, Ont.

ANS.—There are so many causes for lameness that it is impossible to foretell always what the result will be.

It may arise from pressure of a tumour which will be fatal, or it may be caused by cold and so, easily cured.

- (1) Separate the bird so that he may be comfortable and with soft earth, sawdust, straw, etc. to sit upon.
- (2) Feed more soft food than usual.
- (3) After giving one Compound Cathartic pill, give the following daily.
- (4) Of Parrish's chemical food ½ a teaspoonful with 6 drops of Tincture of Nux Vomica, best given in a no. 00 Gelatine Capsule. Report results, please.

Question. 1. Please tell me what care and food must be given to hens and roosters when they are moulting. Age of hens 2 years and age of roosters 3.

2. I have a black-red game cock he has been fighting with some other roosters and his face is all sore, especially his eye and mouth, his eye is closed. What shall I put on it. His age is 3 years.

3. Do you think that by breeding a black-red Game rooster with a blue Game hen I could raise some black breasted Game chickens.

F. H. FLINT.

Toronto.

1. More than usual care in all respects. Beware of draughts cold rains and other sources of chill. Feed more stimulating food than usual as spiced and scalded meals, meat, etc. Do not, however, omit green food if the fowls are confined.

Put a little tincture of iron in the drinking water or that with which you mix the soft food. If in the drinking water, just enough to tinge it faintly to be tasted.

House in the birds at nights so that while free from draughts there shall be abundant ventilation.

2. *Treatment of a wounded cock.* If the brain has not been injured or the bird's strength been too largely expended he will recover. Feed him extra well in spiced food and also meat for a couple of days.

If his eye inflames bathe it well twice a day with hot water, the warmest you can bear on your own cheek.