

Under side of the fore wings silky, fuscous and stained with vinous along the outer part of the costa. Under side of the hind wings lighter than above. Under side of the body, middle and hind legs, white and silky. The fore legs are pale fuscous.

This beautiful little species was collected in Florida and sent to me by Rev. Geo. D. Hulst.

CORRESPONDENCE.

XYLORYCTES SATYRUS.

Dear Sir: In answer to Dr. H. A. Hagen's query in the ENTOMOLOGIST for Dec., 1884, I have to say that some four years ago a boy brought me a living specimen, a fine male, of *Xyloryctes satyrus*, which he had taken in Bleeker's Woods, just outside of the city limits.

JAMES T. BELL, Sc. D.

Belleville, Ont., Feb. 17th, 1885.

Dear Sir: Dr. Hagen asks in Dec. No. (Vol. xvi., p. 239) whether this beetle occurs further north than Pa. I find it recorded by Zesch and Reincke in their list of species captured within fifteen miles of Buffalo; by Howard and Schwarz in list of Coleoptera of lower peninsula of Michigan, presumably from Detroit; by Pettit, as captured at Grimsby, Ont., and by Prof. Bell, as taken near Belleville, Ont. It also occurs here—and this is probably the most northerly record for it—but is apparently rare. The only specimen I possess is a ♂ which was brought to me alive by a friend, but I have several times met with the elytra and other fragments of dead specimens in or under decayed logs. It is of course noticeable as being our largest representative of the Scarabæidæ, which in these colder regions only muster about forty species.

W. HAGUE HARRINGTON.

Ottawa, 25th Feb., 1885.

Dear Sir: I am able to add to the kind answers to my query by Messrs. G. Dimmock and S. Henshaw, two more. Prof. Chas. V. Riley writes me—that August 18th, 1871, he found larvæ, pupæ and fresh beetles of *X. satyrus* quite common at Ridgewood, N. J., under old leaves in the