

Two of these were found in my garden hid among some weeds, June 29, 1880. After putting them in the rearing box one died, the other changed to a chrysalis July 5th, and produced the imago July 18th.

*Prodenia lineatella* Harvey.

Length 1.35 inches. Shape of body cylindrical, very much like the larvæ of *Agrotis*. Dorsal, subdorsal and substigmatal lines, and a line in the middle of subdorsal space, pinkish lilac. In the dorsal space there are, first a series of oval dark brownish drab spots, the broadest part occupying the centre of each joint, the several spots connecting with each other at the union of the joints. In width they extend two thirds of the distance from the dorsal to the subdorsal lines. On the subdorsal rests a series of either triangular or somewhat oval velvety black spots, one to each joint on each side of the body. These spots extend inward half way to the dorsal line, thus encroaching somewhat upon the drab spots. At the base of each black spot, next the subdorsal line, is a narrow, bright yellow, semi-elliptical spot. The rest of the dorsal space is gray irregularly striped with fine white. The upper half of subdorsal space is pinkish gray irregularly striped with fine black; the lower half black, similarly striped with fine light lines. Substigmatal space and venter carneous gray, the first spotted with white. The head and cervical shield are black, the inverted Y white.

The single larva from which the above description was taken was found in my garden August 20th, 1880, in a bed of salsify. While in confinement it ate readily of the leaves of salsify, peach and raspberry. August 25th it disappeared for pupation, and the imago emerged September 11th.

*Leucania pseudargyria* Guen.

Length one inch. General color fleshy brown, sprinkled over with dark brown. Dorsal line narrow, of general ground color; this and the subdorsal and substigmatal distinguished by not being sprinkled with brown. The dorsal space dark brown, composed of dark brown dots finely sprinkled over the surface. About midway this is partially separated as though forming two dark lines. There is a slight massing of these brown dots from the posterior part of the joint, near the dorsal line, outward forming an indistinct V. Subdorsal space much like the dorsal only lighter. Substigmatal space with few dots, a brown patch at the