

PÆDERUS, <i>Grav.</i>	OXYTELUS, <i>Grav.</i>	TRIGONODEMUS, <i>Lec.</i>
Littorarius, <i>Grav.</i>	Sculptus, <i>Grav.</i>	Striatus, <i>Lec.</i>
STENUS, <i>Latr.</i>	TROGOPHILÆUS, <i>Mann.</i>	ANTHOBIUM, <i>Steph.</i>
*Colon, <i>Say.</i>	*Morio, <i>Er.</i>	Protectum, <i>Lec.</i>
Juno, <i>Fab.</i>	ANTHOPHAGUS, <i>Grav.</i>	PROTEINUS, <i>Latr.</i>
OXYPORUS, <i>Fab.</i>	*Cæsus, <i>Er.</i>	Parvulus, <i>Lec.</i>
Rufipennis, <i>Lec.</i>	LESTEA, <i>Latr.</i>	MEGARTHURUS, <i>Steph.</i>
*Femoralis, <i>Grav.</i>	*Biguttula, <i>Lec.</i>	*Americanus, <i>Sachse.</i>
Lateralis, <i>Grav.</i>	ACIDOTA, <i>Steph.</i>	Angulicollis, <i>Makl.</i>
BLEDIUS, <i>Steph.</i>	*Subcarinata, <i>Er.</i>	OLISTHÆRUS, <i>Er.</i>
Fumatus, <i>Lec.</i>	*Patruelis, <i>Lec.</i>	Nitidus, <i>Lec.</i>
PLATYSTETHUS, <i>Mann.</i>	OLOPHRUM, <i>Er.</i>	GLYPTOMA, <i>Er.</i>
Americanus, <i>Er.</i>	*Emarginatum, <i>Er.</i>	*Costale, <i>Er.</i>
HISTERIDÆ.		
HOLOLEPTA, <i>Payk.</i>	Coarctatus, <i>Lec.</i>	*Deletus, <i>Lec.</i>
Fossularis, <i>Say.</i>	*Cylindricus, <i>Payk.</i>	TERETRIUS, <i>Er.</i>
HISTER, <i>Linn.</i>	Gracilis, <i>Lec.</i>	*Americanus, <i>Lec.</i>
Interruptus, <i>Beauv.</i>	PHELISTER, <i>Mars.</i>	PLEGADERUS, <i>Er.</i>
Merdarius, <i>Payk.</i>	*Subrotundus, <i>Mars.</i>	Transversus, <i>Say.</i>
*Harrisii, <i>Kirby.</i>	TRIBALUS, <i>Er.</i>	ACRITUS, <i>Lec.</i>
Fœdatus, <i>Lec.</i>	*Americanus, <i>Lec.</i>	*Exiguus, <i>Lec.</i>
*Cognatus, <i>Lec.</i>	PAROMALUS, <i>Er.</i>	SCAPHIDIIDÆ.
Marginicollis, <i>Lec.</i>	—— ?	SCAPHIDIUM, <i>Oliv.</i>
Depurator, <i>Say.</i>	SAPRINUS, <i>Leach.</i>	4 Pustulatum, <i>Say.</i>
Abbreviatus, <i>Fab.</i>	Distinguendus, <i>Mars.</i>	Piceum, <i>Mels.</i>
*Sedecim striatus, <i>Say.</i>	*Pennsylvanicus, <i>Er.</i>	SCAPHISOMA, <i>Leach.</i>
Americanus, <i>Payk.</i>	Conformis, <i>Lec.</i>	Convexum, <i>Say.</i>
*Carolinus, <i>Payk.</i>	Assimilis, <i>Er.</i>	TOXIDIUM, <i>Lec.</i>
LeContei, <i>Mars.</i>	*Scrupularis, <i>Lec.</i>	Gammaroides, <i>Lec.</i>
Parallelus, <i>Say.</i>	Fraternus, <i>Lec.</i>	

MISCELLANEOUS NOTES.

LARVA OF *HELIOTHIS ARMIGERA*.—At a recent meeting of the Entomological Society of London, Eng., "Mr. J. Jenner Weir exhibited two specimens of *Heliothis armigera*, Linn. (*H. umbrosus*, Grote), bred from larvæ which fed in tomatoes. An importation of tomatoes from Spain or Portugal had been greatly damaged by a number of green larvæ, with black lines and spots, which fed in the fruit, where there was apparently juice enough to drown them, and which ultimately produced the moths exhibited." This

* Species marked with an asterisk have not been before included in the list of Canadian Coleoptera.