D. cyaneo-maculata, nov. sp. (Jac. in litt.).

Head and thorax yellow; antennæ, except the extreme base, black-Thorax bifoveate and depressed; scutel yellow; elytra yellow, slightly dilated at the rear thickly, coarsely and subscriately punctured, with a large cyaneous blue spot occupying the whole apical third, leaving the extreme margin narrowly flavous to the tip. Body beneath paler yellow, legs yellow, with tibiæ, tarsi and upper surface of apex of femora black-Length, 5½ mm.

Five examples, Callanga, Peru.

This species has been distributed with the manuscript name cyaneo-maculata Jac. Of what purport to be three co-types sent me by Messrs. Staudinger & Bang-Haas, two belong to a different genus. The antennæ attain the apical spot; joint 3 a trifle shorter than 4. The thorax is finely but obviously punctulate, while the elytra show here and there fragments of smooth lines between the punctures. In one of the specimens the apical spot includes the side margin, in the typical form it does not.

D. cyaneo-plagiata, nov. sp.

Head yellow; mouth-parts dark; antennæ black, piceous at base, joints 9-11 flavous, with extreme tip of last dark. Thorax flavous, bifoveate, a sublateral dark streak on each side, not attaining either edge; scutel flavous; elytra flavous rufous, dilated behind, coarsely thickly punctured, with a common apical cyaneous blue spot, which does not attain either the side or apex. Body beneath flavous, legs flavous, with apex of femora and tibiæ and tarsi black. Length, 4½ mm.

One example, Peru, green label (Callanga?).

Superficially like cyaneo-maculata, but the light joints of the antennæ and the black streaks of the thorax at once separate it. The elytral spot is smaller; the elytral punctuation is coarse, subseriate in the disk, and becomes obsolete behind and at the sides.

A CORRECTION.

From the description of Gnorimoschema septentrionella, in the December number, p. 422, some words have been left out. The description of the hind wing of the insect should read: Hind wing dark grey—dries with a gloss. Fringe lengthening towards the body to 3½ millimetres at the longest part, light brown.—T. W. F.