

I refer this genus to the Deltoids near *Herminia*. My single specimen in good condition has the fore feet so tufted that I cannot make out satisfactorily the form of the tibiæ and tarsi; they seem to be aborted.

*Pallachira bivittata*, n. s.

Entirely pale ochrey, powdered with fuscous. A broad fuscous stripe below median vein from base to external margin. A second shorter stripe from the extremity of the cell outwardly. Else the entire insect is concolorous. *Expanse* 25 mil. Buffalo, July, coll. auct. The fore wings recall in color those of *Arsilonche Henrici*.

*Agrotis trabalis*, n. s.

♀. Fore tibiæ unarmed; middle and hind pair spinose. Eyes naked. Thorax with a small tuft behind the collar; behind with a divided tuftlet. Abdomen a little flattened, carinated, with a tuft on basal segment. Wings ample; form rather stout. Whitish gray with large stigmata and bright brown contrasting subterminal space. A basal black dash, a second above it on the cell, before the orbicular, which latter is near the t. a. line, inaugurated above it on costa by two black lines rather wide apart, with white included space. Below the t. a. line is twice waved to internal margin. Basal space whitish; basal line indicated. Sub-basal space dark gray. Stigmata concolorous, ringed with black, very large. Claviform incomplete; orbicular a little flattened, ovate; reniform moderately excavate. T. p. line narrow, geminate, regularly and slightly scalloped, with a deeper incision opposite the cell. Subterminal space rather wide, bright brown; s. t. line faint, pale; terminal space rather narrow, dark gray. A dentate black continuous terminal line. Hind wings pale gray fuscous, with pale fringes touched with blackish at extremity of veins; a black terminal line. Beneath pale, powdered with dark scales; an indistinct common outer line; discal lunule filled in and prominent on hind wings, empty on primaries. Abdomen pale; thorax gray, darker shaded on tegulæ. Second palpal joint outwardly black. Front white inferiorly below a frontal black line. The collar has no transverse black line, but is tipped with a darker shade. *Expanse* 42 mil. Mass. (Roland Thaxter).

A second specimen in poor condition, from Montreal (Couper), has the wings more obscure, the brown subterminal space improminent. In the type there is merely a black line inferiorly connecting the stigmata along the median vein. In the Canadian specimen the orbicular and reni-