

Cheeks shiny, finely sparsely punctured, clothed with long sparse white pubescence. Vertex with long sparse hairs, pale and black intermixed, occiput with a short, dense pale fringe.

Prothoracic spine present, short and sharp. Mesothorax with small, round, deep punctures, crowded but distinct anteriorly, along sides and posteriorly well separated, a very few on a good-sized shining disc. Pleura similarly but striately punctured. Scutellum with anterior one-third shining and impunctate, the remainder coarsely, sparsely punctured, and with a median depressed line. Postscutellum densely punctured. Superior face of metathorax fairly well defined, its pits shining, somewhat irregular, longer than broad. Enclosure perfectly funnel-shaped, shining, the bowl convex, with a median and several lateral ridges, the neck concave and perfectly smooth. Surrounding areas shiny, with sparse but very distinct punctures. Pubescence of thorax white, tinged with gray above and sparingly mixed with black on mesothorax and scutellar fringe. Tegulae black, edged with testaceous. Wings subhyaline, nervures dark brown, stigma paler.

Legs stout, black except for apical tarsal joints, which are brownish, clothed with short, dense, white pubescence, that fringing anterior borders of intermediate and posterior tarsi short, stiff and black, that on under surface of posterior femora and tibiae largely black. Basal joint of hind tarsus three times as long as broad, and hind tibiae very stout. Tibial spurs short, dark testaceous, not distinctly pectinate. Claws ferruginous, medially toothed. Anterior coxae with very short blunt spines. Abdomen stout, distinctly subconical, shining black, first segment subimpunctate, or at most indistinctly and scatteringly punctured, second and following segments indistinctly but rather closely punctured, the apical margins perfectly smooth and impunctate under the fasciae, which are broad and pure white on segments 1-5. Basal segment with long, erect, white hairs on the basal truncation, becoming shorter, sparser and subdepressed on the convexity, and forming a short, dense fringe down the sides, uniting with the apical fascia. The following segments have short scattered pale hairs, becoming longer, denser and more bristly in a fringe just before the fasciae, and also intermixed with similar dark ones on the last three segments, especially the apex. Ventral segment 5 deeply emarginate on apical margin. Apical margin of first two segments and base of second segment depressed and constricted.