

Habitat: New Westminster, British Columbia, 1896 (Fletcher); Northwest Territory (Ottolengui).

Two males in good condition, very much alike, yet different. The specimen from Dr. Fletcher came some years ago and was associated with *Momophana Comstocki*, because of the tendency to powder or darken the outer portion of the median space. It was realized that the wing form was somewhat different; but this was not without the range of possible variation, and I had an example from Oregon that I yet believe to be *Comstocki*. The receipt of a fine example from Dr. Ottolengui makes clear a close relationship to *Feralia jocosa*, than which it is a much larger and more intensely coloured form. It is quite likely that the secondaries may vary to almost whitish with more or less blackish powderings.

*Carneades cinereopallidus*, n. sp.—Ground colour a peculiar, very pale ashen gray, more or less powdered with bluish dark gray scales; a little washed locally with luteous. Head concolorous. Collar with a blackish line across the middle, below which is a whitish line or shade which may involve the entire lower half. Tip edged with white scales, then luteous to the black line. Thorax with disc and patagia more or less whitish, the edges of the latter sometimes edged with black: more evidently so in the females. Primaries whitish powdered over the costal region; a luteous shading through the cell and in the s. t. space. A black, geminate basal line is obscurely marked; most obvious on and below the median vein. Median lines practically lost: the t. a. marked by the claviform and a slight difference in shade between basal and median spaces; t. p. a narrow paler line which is somewhat rigidly oblique. S. t. line narrow, whitish, preceded by black scales which may form rather vague sagittate spots. It is obviously dented by whitish rays on veins 3 and 4, which do not, however, quite reach the outer margin. Whitish rays on veins 6 and 7 do not, or only slightly, dent the line. A series of smoky, obscure, terminal lunules. A very pale yellow line at base of fringes. Claviform narrowly outlined by black scales, concolorous, usually with a paler shading above and beyond it. Orbicular oblique, oblong, varying in width, open to the costa, sometimes outlined by black scales, of the palest ground colour or a little whitish. Reniform rather narrow, kidney-shaped, pale yellowish, sometimes contrasting; not outlined except by the rather sharp colour contrast. Secondaries white, with a very narrow smoky edging in the male, a broader, variable outer