

Surgery. Each volume will be complete in itself, and the work will be sold either separately or in sets. Prices per volume: \$3.00 net; half morocco, \$3.75 net. The quantity of matter under review has become so immense that its inclusion in one volume would make a book of unwieldy proportions; but by thus dividing the matter, two volumes will be made of about 600 pages each—an ideal size for convenience of handling and reference. Each volume will be complete in itself, will be paged separately, and will have its own index. This division is made in such a way as to appeal to physicians from a class standpoint, one volume being distinctly medical, and the other distinctly surgical. The apportionment of subjects in each volume is as follows: *Medical*—General Medicine, Pediatrics, Pathology, Nervous and Mental Diseases, Cutaneous Medicine, etc., *Materia Medica*, Physiology, Legal Medicine, Hygiene, Chemistry, etc. *Surgical*—General Surgery, Obstetrics, Gynecology, Orthopedic Surgery, Ophthalmology, Otolaryngology, Laryngology and Rhinology, Anatomy. This arrangement has a two-fold advantage. To the physician who uses the entire book, it offers an increased amount of matter in the most convenient form for easy consultation, and without any increase in price; while the man who wants either the medical or the surgical section alone secures the complete consideration of his branch without the necessity of purchasing matter for which he has no use. All the features that have proved so valuable heretofore will be retained in the two volumes, and the American Year-Book will continue the only adequate interpreter of current medical literature. W. B. Saunders, 925 Walnut Street, Philadelphia.

---

---

### Miscellaneous

---

THE TREATMENT OF POST-PARTUM HEMORRHAGE.—Dr. J. Z. Currie (*Boston Medical and Surgical Journal*, November 16th) summarizes an interesting article on this subject as follows: (1) A knowledge of the source of hemorrhage is necessary to insure intelligent action. (2) All rents, when easy of access, should be repaired at once. (3) If the body of the uterus is contracted and bleeding is excessive, and in all cases of hemorrhage following placenta previa, the general cavity should be tamponed at once. (4) If this is not successful, or if the hemorrhage is constant and not excessive, the bleeding vessels should be secured if possible, and the injury repaired. (5) If atony exists and hemorrhage is not excessive, external and bimanual compression of the uterus should