

*ICHNEUMON OCCIDENTALIS*, n. sp.

Female—Length, 11 mm. Rufo-ferruginous. Head rufous, subtuberculate beneath antennæ, face sparsely punctured; antennæ black, thickened towards apex, with a white annulus, scape rufous. Thorax mostly black; the mesonotum, scutellum and spot at base of metathorax rufous, legs entirely rufous, except apical joint of tarsi, which is brownish. Abdomen rufous; postpetiole and two following segments closely opaquely punctured, remaining segments shining; gastrocœli oblique, linear; a strong transverse depression near apex of the same segment.

Described from one ♀ specimen taken at Victoria, Vancouver Island, in Nov., 1890, by Mr. Taylor.

*PLATYLABUS PACIFICUS*, n. sp.

Female—Length, 9 mm. Black, with rufous abdomen and legs. Head small, closely punctured; palpi reddish, antennæ entirely black, slightly stouter beyond the middle. Thorax closely punctured, the pleuræ more coarsely; metathorax rugose, excavated behind; carinæ indistinct, angles subspinose; legs rufous, coxæ and trochanters black; wings subhyaline, stigma and nervures reddish, areolet pentagonal. Abdomen rufous, polished, except postpetiole, which is aciculated, with two dorsal carinæ not reaching the apex; ovipositor slightly exerted.

Described from one ♀ specimen from Vancouver Island (Taylor).

*CENTETERUS CANADENSIS*, n. sp.

Female—Length, 5 mm. Black; legs and band on abdomen rufous. Head large, vertex and cheeks polished, face punctulate; mandibles rufous, palpi pale; antennæ black, scape oval, red beneath, joints 3-5 subequal, about twice the length of the remaining joints, which are about as broad as long. Thorax black, shining, the mesonotum and pleuræ with sparse faint punctures, the metathorax distinctly areolated, the angles spiniform; legs rufous, the posterior coxæ, tibiæ at apex and tarsi piceous. Abdomen polished, first and apical segments black; two, three and base of four rufous.

Described from 3 ♀ specimens from Victoria, V. I. (Taylor.)

*HERPESTOMUS FLAVICOXÆ*, n. sp.

Male—Length, 6 mm.—Black, with rufous legs. Head black, front rounded, sparsely punctate; clypeus, mandibles and palpi yellow; antennæ stout, black above, piceous beneath, scape yellow. Thorax black,