

Vertex but little rounded above, the apex slightly rounder than in *Kirkaldyi*. Second joint of antennæ very large, consisting of a long, flat plate thickest on the margins and studded with fine knobs; elytra very large, venation as in *Burnetii* nearly, the outer branch of the median nervure straight, the cross nervures at the apices of the elytra in a straight line; costal appendage as long as the second joint of antennæ, strap-shaped towards apex, the posterior margin nearly straight, anterior margin sloping off to the base of the costa; the whole appendix curved back across the corium, with the apex on the claval suture.

Colour: pale straw; eyes black; elytra milky at base, a fuscous band at one-third the distance from base, beyond this subhyaline, with the nervures faintly brown as far as the apical nervures. Posterior margin of appendage, and sometimes a spot near the outer corner of scutellum, fuscous.

Described from two females collected at Albion, N. Y., by E. P. Van Duzee.

*Patara Vanduzeei*, n. sp.—Form and general appearance of *guttata*, but with a smaller front and different venation; brownish purple, with a light line on vertex and pronotum, and light dots around the apex of elytra. Length, 4.75 mm.

Vertex and front together semicircular, about equally margining the eye all around as seen from side, front rising abruptly from clypeus, compressed, the margin slightly thickened; vertex expanded posteriorly; pronotum slightly carinate, broad and nearly parallel margined; venation closely resembling *guttata*, but with four cells between the postcostal and the median before the apical cell. There is a reflexed veinlet from the median into the anal area, and two reflexed veinlets from the outer branch of the mediastinal to the costal. This branch is interrupted just before the apex, making it appear as if the reflexed vein was the end of the nervure.

Colour: vertex white, front and antennæ brownish testaceous; pronotum brownish fuscous, with a broad, median, light stripe; scutellum bright testaceous, sometimes with a pale stripe; elytra brownish purple, the tuberculate nervure of clavus white, a light spot on the apex of each apical nervure and a broad one on the inner reflexed one. Nervures bright testaceous, the three cross nervures before the apical cells fuscous.

Described from three females in the collection of E. P. Van Duzee, from Gowanda, N. Y. The adding of this name to the list is peculiarly