

Sanatoria for the reception of phthisical (paying) patients, II. 72-77

Scarlet fever, II. 636-49

Aconite in, II. 632
Albuminuria in, II. 640
Alcohol in, II. 646
Complications and sequelæ of, II. 647-49
Diaphoretics in, II. 642
Diet in, II. 643
Glandular enlargements in, II. 648
Indications for treatment of, II. 641-47
Isolation and disinfection of patient in, II. 638
Mild forms of, II. 641-44
Notification of, II. 638
Otitis media in, II. 647
Prophylactic measures, II. 636
Quinine in, II. 645, 647
Serumtherapy in, II. 641
Severe forms of, II. 644
Sprays, washes, and gargles in, II. 640
Stimulants in, II. 646
Tenacity of poison of, II. 637
See also Additional Formulæ, II. 667-68

Schott (or Nauheim) treatment of cardiac diseases, I. 419-23, 428-29

Sciatica. See Neuralgia of the sciatic nerve, II. 329-33

Sclerosis, Amyotrophic lateral, II. 307-8

Scurvy or Scorbutus, II. 548-55

Causes of, II. 549-50
Constipation in, II. 544
Curative measure in, II. 552
Diarrhœa in, II. 554
Importance of dietary in, II. 553
Indications for treatment of, II. 552
Infantile, II. 554
Mouth washes in, II. 553
Prophylactic indication in, II. 552
Symptoms of, II. 550-52
Tonics in, II. 553
Warm baths and massage in, II. 554
See also Additional Formulæ, II. 556

Sea-sickness, I. 149-52

Abdominal belt in, I. 152
Amyl nitrite in, I. 152
Atropine in, I. 151
Brandy in, I. 151
Bromide of ammonium in, I. 150, 152
Chloral in, I. 151

Sea-sickness (continued)

Chlorobrom in, I. 152
Chloroform in, I. 150
Cocaine in, I. 150, 151
Enemata in, I. 151
Ice-bag to spine in, I. 152
Iced champagne in, I. 151
Indications for relief of, I. 149, 151
Iodine tincture in, I. 152
Morphine in, I. 151
Oxalate of cerium in, I. 152
Resorcin in, I. 152
See also Additional Formulæ, I. 165, 166

Serum treatment in cerebro-spinal fever, II. 775, 778

in diphtheria, II. 591, 596-601
in dysentery, I. 274
in endocarditis, malignant, I. 359, 360
in erysipelas, II. 664-65
in exophthalmic goitre, I. 530-31
in hay fever, I. 627
in pernicious anemia, I. 508
in phthisis, II. 22, 23
in pneumonia, acute lobar, I. 643
in rheumatism, acute, II. 567
in scarlet fever, II. 641
in tetanus, II. 748
in typhoid fever, II. 676

Shaking palsy. See Paralysis agitans, II. 387-90

Shingles, II. 333-34

Small-pox, II. 620-36

Antiseptic in, II. 633, 635
Chloral and sodium bromide in, II. 632
Cinchona bark in, II. 633
Complications in, II. 635
Confluent cases of, II. 633
Diaphoretics in, II. 628
Diet in, II. 629, 632, 637
Dover's powder in, II. 629
Eruptive stage of, II. 629-32
Hæmorrhagic or malignant, II. 632

Incubation of, II. 626
Indications for treatment of, II. 626

Initial stage of, II. 628
Prophylactic measures, II. 620-26

Quinine in, II. 630
Red light in, II. 630
Sprays, washes, and ointments in, II. 630-35
Stage of regression, II. 635
Stage of suppuration, II. 632
Vaccination in, II. 624-26
Varioloid, II. 624
See also Additional Formulæ II. 666-67