

## LIST OF FIGURES (continued)

<u>Figure Number</u>		<u>Page Number</u>
3-19	Distribution of lakes sampled in Ontario Ministry of the Environment 1981 and 1982 surveys.	3-62
3-20	Mean summer epilimnetic alkalinity, mean summer epilimnetic pH and minimum surface water pH in spring for 16 lakes in Muskoka-Haliburton (1976-1980).	3-63
3-21	Minimum pH values of 57 headwater streams in Muskoka-Haliburton, 1976-80.	3-64
3-22	Calcite saturation indices for 181 lakes in southern Quebec, summer 1980.	3-66
3-23	pH Values for Quebec lakes, summer 1980.	3-67
3-24	$\text{HCO}_3^-/\text{SO}_4^{2-}$ values for Quebec lakes, summer 1980.	3-70
3-25	Distribution of calcite saturation index values for New Brunswick, Prince Edward Island and Nova Scotia.	3-71
3-26	Distribution of calcite saturation index values for Newfoundland.	3-72
3-27	Distribution of calcite saturation index values for Labrador.	3-73
3-28	Distribution of surface water alkalinities for the United States.	See map folio
3-29	Surface water alkalinity of New England states.	3-75
3-30	Annual changes in median pH and mean discharge-weighted excess $\text{SO}_4^{2-}$ for the St. Mary's and Medway Rivers, Nova Scotia, and the Isle aux Morts and Rocky Rivers, Newfoundland.	3-80
3-31	Geographic distribution of pH levels measured in Adirondack lakes higher than 610 metres elevation, June 24-27, 1975.	3-84
3-32	Frequency distribution of pH and fish population status for 40 high elevation lakes surveyed in the 1930s and again in 1975.	3-85