

KENTUCKY BUTTERFLIES.

BY HATTIE H. WARNER, LEXINGTON, KENTUCKY.

The following list has for its foundation a collection belonging to the Experiment Station of the Kentucky State College. The diurnal Lepidoptera have not been made the object of special collecting by the Station entomologist, and doubtless a good many species will be added in the future:—

1. *Papilio ajax*, Linn. Common in all wooded regions throughout the State. High Bridge, Nortonville, Clay's Ferry, etc.
2. *Papilio philenor*, Linn. Frequent throughout the State. Bowling Green, Nortonville, Lexington, etc.
3. *Papilio asterias*, Fab. Frequent everywhere in the State.
4. *Papilio troilus*, Linn. Common locally. Specimens from Glasgow, Clay's Ferry, Fulton, etc.
5. *Papilio turnus*, Linn. Very common everywhere. Specimens from Providence, Lexington, Nortonville, Clay's Ferry, etc.
6. *Papilio cresphontes*, Cram. Rather rare. Lexington, Bowling Green.
7. *Pieris protodice*, Boisd. and Lec. Common some seasons, generally rare. Lexington.
8. *Pieris rapæ*, Linn. Very common everywhere.
9. *Nathalis iole*, Boisd. Very rare.
10. *Callidryas eubule*, Linn. Rather common locally in Western Kentucky.
11. *Colias casonia*, Stoll. Lexington. Not common.
12. *Colias philodice*, Godt. The most abundant butterfly of Kentucky. Occurs in large swarms about damp sand in the latter part of summer.
13. *Terias nicippe*, Cram. Common everywhere in lowlands. Lexington, Nicholasville, Bowling Green, High Bridge, etc.
14. *Terias lisa*, Boisd. and Lec. Found only occasionally. Specimens from Glasgow.
15. *Danais archippus*, Fab. Common everywhere.
16. *Argynnis diana*, Cram. Very rare. A specimen was collected by Prof. H. Garman, at East Cairo, several years ago.
17. *Argynnis cybele*, Fab. Common about thistles in summer.
18. *Euptoicta claudia*, Cram. Common in fall all over the State. Lexington, Bowling Green, East Hickman.