

brooded. From the fact that the exuding pitch offers so much protection to the insect, it is scarcely likely that any remedy would reach it. The knife seems to be the only resource. A small four-winged parasite attacks the borer in the larval state, the chrysalis being often found filled with the cocoons of this useful friend.

COMSTOCK'S RETINIA—*Retinia Comstockiana*, Fernald.

This insect as yet has only been observed on the pitch pine (*Pinus rigida*), but as it is probable that it will sooner or later be found to attack the white pine it will be briefly noticed.

The perfect insect is a small moth of a light grey colour, varied with darker shades of rusty brown, which measures, when its wings are spread, nearly three-quarters of an inch across. The eggs are laid on the terminal shoots of the trees, and the larvæ are found in the early part of the summer boring into the twigs and small branches, causing an exudation of resin, and sometimes girdling them. The larvæ, when full-grown, are nearly half an inch long, of a yellowish colour, with a brown head, a patch of the same colour on the next segment, and a few polished brown dots on each ring, and from every one of these there arises a single hair. The larva changes to a chrysalis within the burrow from which eventually the moth escapes. In fig. 28 we have the insect represented in its several stages, also a small branch of an affected tree and a section of one of the bored twigs.

THE PINE LEAF-MINER—*Gelechia pinifoliella* (Comstock).

This insect was first described by Prof. Comstock in his report to the Department of Agriculture, Washington, for the year 1879. It has been found mining the leaves of different species of pine in many parts of the United States, and although as yet unrecorded in Canada, will very probably be found here. The larva is very minute, in the figure it is much magnified; the line below indicates the natural size. It lives within the leaf on the soft tissues, and its presence is soon indicated by the change in colour which takes place in the part affected; it becomes brown, and on examining the leaf, or that portion of it which is discoloured, it is found to be entirely eaten out, and to contain, if in season, the insect, either in the larval or pupal condition. In fig. 29 the work of this tiny insect on the leaves is shown. The moth, chrysalis, and larvæ are all represented, but much magnified.

The larva, when full-grown, is about one-fifth of an inch long, of a pale brown colour, with a black head and a black patch on the upper part of the next segment. It is also sparingly covered with short, fine hairs. The change to a chrysalis takes place within the mined leaf, and in summer the moth escapes in about a fortnight.

The perfect insect, when its wings are spread, measures about three-eighths of an inch across; it is of a brownish-yellow colour, dotted with fuscous scales. The fore-wings are crossed by three white lines, as shown in the figure; the hind-wings are pale grey,




Fig. 28.

and bot  
the year

In ad  
are known  
Unfortun  
towards li  
deter us fr  
much mor  
or orname  
of Paris g  
water. Ba  
broom; su  
a lodgemen  
protected;  
areas of fo  
insectivoro  
insects, whi  
these friend  
of trees an

This fo  
New York,  
damage. T  
domiciled its  
to be on the  
economic en  
During  
York State I  
Smith, of R