

9. **Limnophilus ornatus** Banks (?).

Limnophilus ornatus (?) Ulmer, Cat. Coll. Zool. Selys, VI (1) p. 20, figs. 30, 31, taf. I, fig. 6 (1907); Ulmer, Deutch. Ent. Zeit., ip. 341 (1908), nec Banks.

This species is unknown to me. Ulmer recorded two female specimens from Hokkaido ("Yerse"), saying, "Die beiden Stücke stimmen gut, wie das auch meiner Figur auf Tafel I zeigt, mit der Beschreibung von Banks überein; die Appendices præanales eines amerikanischen Exemplares, das mir von C. Betten gesandt wurde (det. N. Banks) sind allerdings an der Basis etwas breiter und schwach medianwärts statt lateralwärts gekrümmt; . . . "

In any case it is very interesting to know that the same or a very closely allied species occur in such widely separated localities as North America and Hokkaido.

Genus *Nothopsyche* Banks.

10. **Nothopsyche pallipes** Banks.

Nothopsyche pallipes Banks, Proc. Ent. Soc. Wash., VII, p. 107, pl. III, fig. 1 (1906); Ulmer, Cat. Coll. Zool. Selys, VI (1) p. 29, figs. 48, 49 (1907); Ulmer, Deutch. Ent. Zeit., p. 341 (1908).

Habitat—Hondo (Gifu, and Numata, Prov. Kozuke).

Time of appearance—October.

11. **Nothopsyche ruficollis** (Ulmer).

Chilostigma ruficolle Ulmer, Stett. Ent. Zeit., Jg. 66, p. 14, taf. I, figs. 12, 13 (1905).

Nothopsyche ruficolle Banks, Proc. Ent. Soc. Wash., VII, p. 107 (1906).

Nothopsyche ruficollis Ulmer, Cat. Coll. Zool. Selys, VI (1) p. 29, figs. 46, 47 (1907); Ulmer, Deutch. Ent. Zeit., p. 342 (1908).

Habitat—Hondo (Tokyo, Gifu, Numata). Shikoku (Uwajima, Matsayama). Kiushin (Yanagawa, Prov. Chikugo).

Time of appearance—August to November.

12. **Nothopsyche longicornis**, sp. nov.

Head fuscous black; frons with long piceous hairs; maxillary palpus greyish black; antenna black, much longer than fore-wing.