

BROMIDIA IN SLEEPLESSNESS.

For nervousness, sleeplessness, and sexual excitement, characterized by erections or even chordee, various authorities vary in their recommendations. Ringer recommends the use of aconite and camphor. Bartholow and Phillips both advise the administration of lupulin. The value of hyoscyamus has been appreciated by many medical men for a long time, and is quite valuable. Bromidia is to be highly recommended, since it consists of chloral, bromide, hyoscyamus and cannabis indica, and acts as a somnifacient, spinal sedative, and hypnotic. The dose is a drachm to two drachms an hour before bed time.—*American Journal Dermatology.*

TREATMENT OF UTEROVAGINAL CATARRH.

Fifteen months ago Mrs. X. came to me for treatment, giving the following history: Six years previous she had a miscarriage, since which she had been troubled with a profuse leucorrhœa of a very foul odor. At her menstrual period she suffered greatly and flowed excessively. On examination the cervix was found to be nearly four times its normal size and so badly eroded as to have every appearance of a cancer and had been mistaken for such by one physician. The uterus was soft and boggy and very much enlarged. She had been to the hospital on two occasions and each time had been curetted, but this seemed only to aggravate the general condition. For over a year I treated her with every means at hand, but to no purpose. I was making preparation for an operation, which would have meant the removal of the uterus, when my attention was drawn to Glyco-Thymoline and I determined to give it a thorough trial before operative measures were to be further indicated. An intra-uterine douche of Glyco-Thymoline in 25 per cent. hot solution was administered and lamb's wool tampons saturated with Glyco-Thymoline pure were used. She began to improve from the first application. The leucorrhœa became less and the odor disappeared entirely. The cervix took on a healthy look. The uterus decreased in size and became firm; in fact she is now nearly well after nine weeks' treatment with Glyco-Thymoline.

SOME OF THE INDICATIONS FOR SANMETTO.

Vesical irritation and atony; enuresis due to atony; incontinence of urine in children due to a weak bladder; dribbling of the urine in the aged, not due to paralysis or growths; urine expelled upon exertion, as coughing; cystitis; catarrhal discharges from bladder or genitalia of male or female; seminal emissions; prostatitis, enlarged prostate and pre-senility.