

about as long as tibia, but little broader, broadest near base, fingers not near as long as hand, quite strongly curved, and darker than hand, all parts with fine simple hairs, about one-half the width of the joints, except some at the base of the fingers. Abdomen moderately broad, flat, the setæ not distinctly divided, but apparently some of the posterior ones are divided.

Length, 2.5 mm.

Punta Gorda, Florida (Mrs. A. T. Slosson).

*Ideobisium tibiale*, n. sp.—Cephalothorax and palpi red-brown, rest of body and legs pale. Cephalothorax a little longer than broad, rather broader behind than in front, surface smooth, with a few long bristles, anterior margin rather prominent in the middle, two eyes each side, about one-half their diameter apart; mandibles large, with a few long hairs, stylet slender, trifid at tip. Palpi quite long; trochanter nearly twice as long as broad, sides subparallel; femur plainly longer than the cephalothorax, enlarging from base to tip; tibia distinctly shorter than femur, long pedicellate, but one and a-fourth broader than femur, inner side but little swollen; hand as long as femur, not twice as broad as tibia, but little swollen, mostly on inner side, fingers plainly shorter than hand, stout and but little curved; all with long simple hairs. Abdomen not very elongate, scuta smooth, with a few simple hairs; femora i and ii divided, basal part longer than apical part; femora iii and iv showing trochantins distinct.

Length, 4.2 mm.

From Florissant, Colo., 8,000 feet.

*Ideobisium magnum*, n. sp.—Cephalothorax and palpi dark red-brown, dorsal scutæ brown, rest of body and legs pale. Cephalothorax smooth, about as broad as long, a rounded tubercle in middle of front margin, eyes large, less than one-half their diameter apart, a few pale spots each side on the posterior part of the cephalothorax; mandibles large, with a few long hairs, stylet slender, with an outer branch toward tip, the latter bifid. Trochanter of palpus concave behind; femur subpedicellate, hardly as long as the cephalothorax plus the mandibles, as broad at middle as at tip; tibia plainly shorter than femur, strongly pedicellate, inner side suddenly swollen and then straight; hand large, as long as tibia, and about twice as broad, tapering to the stout fingers, which are plainly a little shorter than the hand; all parts with fine simple hairs.