

that preceded it. Now her functions are regular, and I know no reason why she may not become pregnant.

Case 2.—Mrs. S. complained of a continuous, dull, dragging pain, situated in the region of the iliac fossa of the right side. Menstruation irregular, scanty, fetid. Married six years; had never been pregnant. Excessive leucorrhea, though otherwise she was perfectly normal. Her weight was 140 pounds. Her condition, and the suffering, both physical and mental, which it occasioned her, was rapidly undermining her health. She was becoming emaciated, appetite of no consequence, general weakness. She considered her condition "hopeless." Cardiac weakness, of which she was a victim, contra-indicated curettement, which usually cures "whites" and allows the formation of a healthy lining membrane. Ergoapial (Smith) was prescribed for her, one capsule three times a day. In conjunction with this I gave tonic medicines. After six weeks' use of this remedy, the woman said she was "feeling so good" that she did not need any further treatment. She had increased in weight, and her appetite had become all she could wish. The menstrual flow was increased, and now, five periods having elapsed from the time treatment was instituted, her monthly flow has failed to appear. She does not expect its return for some time—supposing herself pregnant.

Case 3.—Miss S. suffered severe pain each month, beginning a day before the flow came on. The flow was a thick, clotted mass, consisting of membrane and the menstrual blood matted together. She had suffered from puberty, and the suffering became more intense as the years passed on. She was 19 years of age, stout, of healthy parentage. Admitted to Lincoln Hospital, January 15th, 1902. She declined an operation. I afterwards prescribed Ergoapial (Smith) and have continued it for one month. Her next menstruation was free and easy; painless and regular. I doubt not that keeping up this treatment up to another period she will be entirely rid of the hitherto troublesome condition.

Case 4.—Miss W., tubercular history. Menstruation very irregular, sometimes three, sometimes five weeks between periods; very painful; scanty. I prescribed Ergoapial (Smith) one capsule four times a day beginning one week before the menstrual period and continued a week after the period. As a result of this treatment the patient feels a great deal better in her general health; her monthly flow has been rendered painless and increased in quantity. Ergoapial has a tonic action upon the muscular fibers of the womb. Its effect is not transitory but lasting. This superior preparation is decidedly tonic.

Case 5.—Mrs. D., a victim of endometritis. Pain continues