

***Satyrus alope*, Fabr.**

(CERCYONIS ALOPE.)

HIPPARCHIA ALOPE.

This species is reported from Fredericton, but appears to be rare in the northern sections of the province. It is very common in the vicinity of St. John. Frequenting the same localities and flying in company with *S. nephele*.

***Satyrus alope nephele*, Scud.**

This butterfly, apparently a hybrid between *alope* and *nephele*, is sometimes taken at St. John.

Family LYCÆNIDÆ.**Sub-family LYCÆNINÆ.*****Thecla angustus*, Kirby.**

(INCISALIA AUGUSTUS.)

Not uncommon, but difficult to capture, owing to its small size, dark color, and the shrubby localities which it frequents. Flies in May.

***Chrysophanus epixanthe*, Boisd Lec.**

(EPIDEMIA EPIXANTHE.)

This little butterfly is not uncommon in two or three localities near St. John, but it appears to be very local in its habits, frequenting the same places (sometimes only a few yards in extent) year after year. Flies in July and possibly into August.

***Chrysophanus hypophlaeas*, Boisd.**

CHRYSOPHANUS AMERICANA, D'Urban.

(HEODES HYPOPHLAEAS.)

Common at St. John, Fredericton, Sussex and Grand Lake, from June to September. This species is no doubt abundant in every part of New Brunswick.

***Lycæna pseudargiolus*, Boisd Lec.**

(CYANIRIS PSEUDARGIOLUS.)

This butterfly is very common throughout the province. The forms *VIO-LACEA*, Edw., being very abundant in May and June, and the form *NEGLECTA*, Edw., less common, in June, July and August.