difference is the rufous violet-colour on the breast, and the want of the long outer tail feathers. The comparing the two figures in the *Planches enluminées*, seems much to favour this opinion. This last may perhaps be a female, or a young bird; for the long tail feathers do not appear till the second year.

Buffon mentions a variety, which came from Goa, and had not the fulvous band on the breast, like the last-mentioned; instead of which, there was a collar of a vinaceous colour bestind the neck, a little below the head.

Coracias orientalis, Lin. Syft. i. p. 159. No 4.

Le Rollier des Indes, Brif. orn. ii. p. 75. pl. 7. f. 2.—Buf. cif. iii. p. 147.—

Pl. enl. 619.

6. ORIENTAL R.

DESCRIPTION.

SIZE of a Jay: length ten inches and a half. The bill is yellowish, broader \* at the base, and more hooked than in any of the genus: the head and hind part of the neck brown: back, rump, scapulars, wing and tail coverts, green brown: throat fine blue; down the shaft of each seather a pale line: the rest of the parts beneath blue green: quills mixed blue and black; on the middle a large pale blue spot: wings longer than in other Rollers: tail even at the end; the two middle seathers green at the base, the rest of the way black; the other seathers are first green, then blue, with black tips: legs yellowish: claws black.

This is a native of the East-Indies.

PLACE.

· Hift. des oif. iii. 147.