

SPRAY CALENDAR

PLANT AND PESTS.	1ST APPLICATION.	2ND APPLICATION.	3RD APPLICATION.	REMARKS.
APPLE.	Earlier before or soon after the leaf-buds burst, preferably the latter. Use A1 or B. For San José Scale prune severely, scrape off loose bark and drench the whole tree, paying special attention to outer twigs.	Just before the blossoms open. Use A2 or D, with 2 or 3 lbs. arsenate of lead to each 40 gals. of the liquid.	Immediately after the blossoms have all, or nearly all, fallen, and before the early ones close. Use A3 or D, with 2 lbs. arsenate of lead to each 40 gals. This is the application for codling moth.	For Scab, a 4th application about 10 days after the 3rd is necessary if June is wet, also an intermediate one between the 2nd and 3rd with A3, <i>without any poison</i>, if the interval, owing to cool damp weather, threatens to be long. Spraying with the weaker A3 early in August is an insurance against sooty fungus and late scab. If Aphids are annually troublesome, delay 1st application till buds begin to burst, then add Black Leaf 40 or nicotine-sulphate 40 per cent. to A1 or B and cover every bud. For Cankers cut out diseased bark, disinfect and cover with white-lead paint free from turpentine. For Blight on young trees keep suckers rubbed off trunk and main branches and cut out promptly any diseased branches or twigs well below the diseased bark. Always disinfect both cuts and tools with corrosive sublimate (1 to 1,000).



Stages for 1st application. Stage for 2nd application. Stage for 3rd application.