

these sources of evidence. Many women are proverbially inaccurate as to dates and in the description of symptoms, and we must ever be on our guard against the designing woman, legitimately or illegitimately pregnant, who wishes to rid herself of the conception, and who hopes that by the use of the sound or other instrument incautiously used by the practitioner, her purpose may be effected. While history, symptoms, and the condition of the breasts are all important, the supreme value in the estimation of the various sources of evidence is to be placed on the bimanual palpation of the uterus. I am in the habit of impressing this on my students. If, with empty bladder and rectum, and everything else favourable in the position of the patient, you cannot easily define the uterine body, so distinctly firm in the nulliparous condition, then suspect pregnancy. It is thus soft in the condition of pregnancy, and comes nearly to the feel of the roof of the vagina and other structures in the pelvis. If the uterus can be defined, the value of the so-called Hegar's sign,—the sudden increase of size above the junction of the body and the cervix—is very great. It is in early pregnancy that mistakes in diagnosis are most frequently made, but I have known not a few in the more advanced stages. Cases are not unknown of all the arrangements having been made for operation for ovariectomy, and the patient meanwhile being delivered of a full term child. This has occurred to men of world-wide reputation, the authors of books and numerous papers on obstetrical and gynaecological subjects. In one instance which occurred to me, ovarian cyst had been diagnosed, and the woman being in great distress from the enormous distension, she had been twice tapped. She travelled over five hundred miles to reach me for operation, all the preliminaries having been arranged. I found her resting on her hands and knees in my waiting room and in that position she had remained during the night in the sleeping car. On examination, I was immediately able, through the cervix, to recognize foetal parts. The case proved to be one of twin pregnancy with hydramnios. The gravid uterus had been tapped and the liquor amnii drawn off. Beyond a doubt the true nature of this case would have been recognized by a careful consideration of history, symptoms, and physical signs, instead of by the mental attitude of taking certain things for granted. Recorded instances are by no means single of operators, when doing hysterectomy for fibroid, being surprised by the discovery of early pregnancy. It is safe to say, from what we know of the very human nature of our profession, that many more have never been recorded. It is doubtless true that operation was the best course in many such cases. The sudden increased activity of growth of fibroids previously unsuspected, in the gravid condition of the uterus, certainly in many such