

Cost of a Medical Education in Great Britain.

According to the *British Medical Journal*, the cost for a student to qualify for the Conjoint Board at a London school is from \$3,250 to \$3,500; or if he proceed to a University degree as much as \$7,000. A perpetual ticket for tuition fees at a London school varies from \$500 to \$750. This of course, does not include the private coaching in different subjects which is commonly found necessary in preparation for examinations. As to the prospect of the newly qualified practitioner, the medical services of the Army and Navy or the Indian service are open to him. If he accept an assistantship with a successful practitioner, he may receive from \$500 to \$750 a year with the prospect of rising to \$1,000 or \$1,250. The *Journal* concludes by stating that the practitioners' services have a definite money value which ought, in the course of years, to give him a fair interest on the capital expended on his education, and, if he be at all successful, by degrees, the return of the capital itself. This is certainly not a prospect to appeal to the imagination of the youth bent on acquiring wealth.

Urotropin in Posterior Urethritis

Gerald Dalton (*The Therapist*) highly commends this remedy in the above named affection. In several of his cases a complete cure was obtained and in none did it fail to produce improvement. He administers 7 grains thrice daily. It is also useful in cystitis and as a uric acid solvent.

W. B. Saunders & Co.'s London Branch.

We are pleased to note that this enterprising publishing house are extending their business by opening a branch in London, England, for the purpose of increasing their business with Great Britain and her colonies. We have no doubt that the same methods that have contributed to the phenomenal success of this company in America, will meet with equal favor across the Atlantic.

Exophthalmic-goitre.

Dr. W. H. Harland (*B. M. J.*, Sept. 1st) reports two cases of exophthalmic goitre with all the characteristic symptoms of the disease, developing suddenly in soldiers after being in action. During the two weeks they were under his observation, no benefit was gotten from treatment. The cases are of interest as confirmatory of the nervous origin of the malady.