THE TREATMENT OF GRANULAR PHARYNGITIS.

M. MOURE (*Presse médicale*; Journal de Bruxelles, June 19th) recommends painting the back of the throat twice weekly with the following:

| Ŗ. | Iodine | 0.25 | grammes (33 grains) |
|----|------------------|--------|---------------------|
| | Potassium iodide | 0.30 | grammes (41 grains) |
| | Laudanum | 3.00 | grammes (45 minims) |
| | Glycerin | 120.02 | grammes (4 ounces) |
| M | | | • |

This may also Le used as a gargle in a strength of a teaspoonful to half a tumblerful or a tumblerful of tepid water.

The author also recommends the use in similar proportions as a gargle, or pure as a local application, of the following :

| B, | Sodium biborate | | | (90 grains) |
|----|--|-------|---------|--|
| | Antipyrine | 4 | grammes | (60 grains) |
| | Tincture of guaiacum Spirit of peppermint | aa 5 | grammes | (75 minims) |
| | Neutral glycerin | . 140 | grammes | (4 ¹ / ₂ ounces) |
| M | | | | |

GLYCOZONE IN PTOMAINE POISONING.

D^{R.} A. RIXA, in the Medical summary, writes as follows regarding some cases of poisoning by fish and oysters: "I prescribed an antespitic intestinal wash, Glycozone, two ounces, hot water, twenty-four ounces, for mornings and evenings. At my evening's call the temperature was 100; pulse 110; respiration, 28. Having had some favorable experience with the internal use of Glycozone in acute gastritis I then prescribed a teaspoonful to be given, diluted with water, every three hours. This treatment was kept up for a week, until all unfavorable symptoms disappeared"

SANMETTO IN PROSTATITIS, ENURESIS, CATARRH OF BLADDER.

「「「「」」」を、「「」」」」と、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、「」」」を、

In prostatitis, enuresis, catarrh of bladder and all diseases of the genito-urinary system, J. T. W. Kerns, M.D., Bellaire, Ohio. has found Sanmetto indispensable to him.

BOVININE IN ENDOMETRITIS.

In this condition much benefit has been derived from the topical and internal employment of bovinine. The uterus is washed out with bovinine two parts and salt solution one part, and the vagina tamponed with pure bovinine. Two teaspoonfuls were given internally every hour in peptonized milk. The local treatment was done twice daty.