

s bracteis-

s oblongo-
floralibus

p. 313.

bsessilibus
bracteisquelanceolatis
que laxis
, genitili-alibus rectia,
limbo subbi-
obæqualibus.ares, loculis
us minutis.

politarii, ter-

o-lineari-
ue viridi-
ato-ovatis
s muticis.earibusve
solitariis,
calycibus
33.

in Linn.

ores duplo
ore erecto
ibus longi-
s. Styles
ad apices
enth.

1. *P. Douglasii*; foliis floralibus bracteisque flores superantibus, staminibus corolla brevioribus, dentibus calycinis inferioribus tubo plus duplo longioribus. *Benth. Lab.* p. 414.

2. *P. multiflora*; foliis floralibus bracteisque corolla brevioribus, staminibus exsertis. *Benth. Lab.* p. 414.

3. *P. parviflora*; foliis floralibus flores superantibus, staminibus corolla brevioribus, dentibus calycinis inferioribus vix tubo longioribus. *Benth. Lab.* p. 414.

1. *Scutellaria* (Sect. *Galericularia*) *tuberosa*; caule humili simplici erecto villosa, foliis petiolatis lato-ovatis obtusis grosse crenatis basi rotundato-truncatis, floralibus oblongis omnibus utrinque villosis, floribus axillaribus oppositis secundis, corollis villosis calyce plus quadruplo longioribus. *Benth. Lab.* p. 442.

1. *Stachys ajugoides*. *Benth. in Linn. v. 6. p. 80.* *Hook. et Arn. supra*, p. 155.

1. *Sphacele calycina*; suffruticosa? ramis molliter villosa, foliis petiolatis ovato-oblongis basi rotundatis subrugosis villosis, racemis subsimplicibus secundis, verticillastris bifloris remotis, calycibus amplis villosissimis, corollis amplis calyces breviter superantibus. *Benth. Lab.* p. 568.

1. *Marrubium vulgare*. *Linn. Sp. Pl. p. 816.*—*M. hamatum*. *Humb. et Kunth, Nov. Gen. et Sp. v. 2. p. 310.*—*M. Germanicum*. *Steud. Nom. Bot. p. 510.*

1. *Trichostemma* (Sect. *Orthopodium*) *lanatum*; foliis linearibus margine revolutis, floribus racemosis lanatis, calycibus subæqualiter 5-fidis, corollæ tubo longe exerto. *Benth. Lab.* p. 659.

2. T. (*Orthopodium*) *lanceolatum*; villosum, foliis lanceolatis, cymis axillaribus, calycibus subæqualiter 5-fidis, corollæ tubo exerto. *Benth. Lab.* p. 659.

ORD. XLV. VERBENACEÆ. Juss.

1. *Verbena lasiostachys*. *Link?*—*Hook. et Arn. supra*, p. 156.

Although a less diffuse specimen than that which we characterized in Captain Beechey's collection, the two are precisely the same species.

ORD. XLVI. PRIMULACEÆ. Vent.

1. *Anagallis arvensis*. L.

This is the common red-flowered variety.

1. *Dodecatheon integrifolium*. Mich.—*Hook. Fl. Bor. Am. v. 2. p. 119.* *Bot. Mag. t. 3622.*—*a.* tubo stamineo flavo. *Hook. Fl. Bor. Am. l. c.*—*y.* tubo stamineo atro-purpureo. *Hook. Flor. Bor. Am. l. c.*

a. Grand Road in the Snake Country. *Tolmie.*—*y.* *Douglas.*—In the *Flor. Bor. Am.* the character of var. *y.* was made to depend on the much greater breadth of the leaves, as well as upon the different colour