

typical dusk-flier. Mr. McGlashan informs us that he always takes his specimens at dusk; the Hymers specimens were also captured at this time of day. Taking all the above points into consideration, we are inclined to think that, if not synonymous with *ganna*, *MacGlashani* is a perfectly valid species.

H. Matthewi H. Edw.

Species described from Brit. Columbia, listed by Dyar as variety of *hyperboreus*. Dr. Barnes possesses only a single rubbed male from Victoria, B. C., which is too poor to make much of. From the description it might be closer to *MacGlashani* than *hyperboreus*. The male alone was described. It behooves our Western friends to furnish good series of this in both sexes.

H. confusus H. Edw.

The type was a single specimen from Alaska. The silver banding appears from the description to be replaced by pale fawn, but we have seen neither the type nor any specimens agreeing with the description. We do not know Dr. Dyar's reasons for placing it as a variety of *hyperboreus*.

H. roseicaput N. & D.

Described from a single male (?) from Cascade Mts., B. C. The description reads rather like that of our female *pulcher* from Mt. Hood, except that this latter species has no blackish subterminal line. Possibly this may be the ♀ of *Matthewi*, described in error as a male; the localities are certainly closely approached.

A NEW SPECIES OF THE SCELIONID GENUS *ACOLOIDES*
HOWARD.

BY A. ARSENE GIRAULT, CHAMPAIGN, ILL.

Superfamily Proctotrypoidea.

Family Scelionidæ.

Subfamily Bæinæ.

Genus *Acoloides* Howard.

1. *Acoloides aureus*, species nova.

Normal position.

Female.—Length, 1 mm. Moderate in size for the genus.

Usual to the genus with the exception of the parapsidal furrows, which are slightly indicated. Winged, the wings short, with moderately

August, 1911