

12 Horses which did not show any improvement by injection of Mallein at intervals of about 30, 60 and 90 days. Although never developing any clinical symptoms.

No	Maximum Temperatures at:				Maximum size of Swelling at :-			
	1st Test	2nd Test	3rd Test	4th Test	1st Test	2nd Test	3rd Test	4th Test
1 Brown gelding, 13 yrs.	105 4-5	105 2-5	105	104 3-5	2x3	4x4	3x4	4x4
2 Bay mare, 10 yrs.	106 2-5	106 2-5	106 1-5		4x6	2x5	6x8	
3 Bay gelding, 8 yrs.	106	105 4-5	105		3x4	5x8	5x6	
4 Roan " 11 yrs.	106 1-5	105 1-5	104 1-5	105 3-5	3x7	4x5	3x5	5x7
5 Chestnut " 14 yrs.	103 1-5	105 2-5	104 3-5	105 2-5	3x6	4x6	5x6	6x6
6 Black " 9 yrs.	105 2-5	105 2-5	105		4x5	3x5	4x5	
7 Bay mare, 9 yrs.	105 4-5	105 1-5	105 3-5		6x7	6x6	6x8	
8 Black gelding, 13 yrs.	105	105 2-5	105 2-5		3x4	2x3	4x7	
9 Brown " 10 yrs.	104	105 4-5	105		6x8	5x5	2x3	
10 Bay mare, 7 yrs.	105 3-5	104 2-5	103 2-5	105 2-5	1x1	3x6	3x4	2x2
11 Bay gelding, 12 yrs.	105 2-5	105 4-5	104 4-5		1x3	2x3	2x3	
12 Chestnut " 8 yrs.	104 4-5	103 2-5	104 1-5	105 2-5	4x6	6x6	2x3	5x6