

and peduncle woolly-tomentose ; rays, numerous, short : achenes hirsutulous ; pappus soft, white, barbellate.

Not rare on the eastern slopes of the Rocky Mountains, at high altitudes, between the International Boundary and Lat. 54°. Distributed from the Herbarium of the Geological Survey of Canada, under numbers 11,606, 14,708 and 19,635.

SENECIO ROBINSII, Oakes.

*S. aureus*. *L. var. lanceolatus*, Oakes ; Macoun, Cat. Can. Plants, vol. 1, p. 265.

A well defined species, easily separable from *S. aureus* and all its varieties. Rare on Cape Breton Island, N. S. The only specimens found by Prof. Macoun in 1898, were at Big Intervale, Margaree, and in woods at Baddeck.

VACCINIUM PENNSYLVANICUM, Lam. var. ANGUSTIFOLIUM, Gray.

Macoun, Cat. Can. Plants, vol. 1, p. 290.

Union Road, Prince Edward Island ; in bogs, Louisburg, Cape Breton Island. (*John Macoun*.) Recent explorations have shown this plant to be widely distributed in Labrador.

PTEROSPORA ANDROMEDA, Nutt.

A new locality for this widely distributed but seldom collected plant is Tête Jaune Cache, headwaters of Fraser River, Rocky Mountains. (*W. Spreadborough*.)

BARTONIA IODANDRA, Robinson.

Holyrood, Newfoundland. (*Robinson & Schrenk*.) Grand Lake, Newfoundland. (*A. Waghorne*.) These records in Botanical Gazette, vol. xxvi, p. 47. In a bog six miles from Half Way House, Cape Breton Island, N. S., Aug. 5th, 1898. Herb. No. 19,857. (*John Macoun*.)

VERBENA STRICTA, Vent.

Along the Grand Trunk Railway embankment at Stamford, Ont., 1898. (*R. Cameron*.) St. David's, Ont., 1898. (*W. Scott*.) New to Canada.