America, north of Mexico. Provancher's *Platymetopius acutus* is the species here identified as *latus*, while his *magdalensis* is the *acutus* of Say. *Platymetopius ornatus*, Baker, has been placed in genus *Dicyphonia* by Dr. Ball.

Deltocephalus Sayi, Fitch.-Apparently common.

Deltocephalus abdominalis, Fabr.—Five examples were taken at Temiskamingue, Quinze Lake and Barrier River. This species may be roughly distinguished from the green examples of debilis by their having the base of the front abruptly black.

Deltocephalus affinis, Baker.—Temiskamingue and Barrier Lake. In compliance with the now generally accepted synonomy of this species, I have applied this name to the species formerly determined by me as Melscheimeri, Fh.

Deltocephalus Melscheimeri, Fitch.—Nigger Point, Quinze Lake. One example. This is a smaller and more slender species than the preceding, and much less abundant.

Deltocephalus inimicus, Say.—Common at all places where Mr. Palmer collected.

Athysanus instabilis, Van D.—Temiskamingue and Quinze Lake. Three examples. These specimens are typical instabilis, which Osborn and Ball identify with the European striatulus, Fall., in which they are very likely correct, but as my material does not agree with any descriptions of that species accessible to me, I prefer for the present to place it under a name of which I am certain, leaving it to future study to settle the synonomy.

Athysanus, sp.—Two examples, representing both sexes, were taken with the preceding. This is a smaller and more slender species, which may be best distinguished by the pale gray elytra with the areoles but obscurely bordered with darker, by the conspicuously white or pale yellow margins to the black abdomen, by its having the pygofers of the female whitish, with their base and the oviduct black, and by the mostly pale legs. This is the species listed as striatulus in my list of the Hemiptera taken by Mr. Palmer at Lake Temagami.

Phtepsius apertus, Van D.—Three examples from about Quinze Lake. An interesting species that finds its home in the far north.

Phlepsius fulvidorsum, Fitch.—Two specimens taken with the last.

Scaphoideus immistus, Say.—Taken at White Rapids and about Quinze Lake.