The type is in the collection of the Union States Biological Survey, the paratypes are in the National Museum and the collection of the author.

Tribe Antochini. Genus Teucholabis Osten Sacken.

Teucholabis lucida, sp. n.

Male-Length 3.5-4.1 mm.; wing 4.2-4.5 mm.

Rostrum and palpi black. Antennæ black. Head black with a thick light gray pubescence.

Pronotal scutum dark brownish black, the caudal margin paler, brown; scutellum pale yellowish white, interrupted medially above by a brownish depression. Mesonotal præscutum shiny black, pilose medially behind, the usual pale interspaces represented only by a small reddish brown area in front, just proximad of the pseudosutural foveæ, and a yellow median patch behind; scutum black shiny, with the median area in front pale yellow and with a short, pale pilosity; scutellum dull yellow, more grayish basally; postnotum black with a gray pile. Pleura light yellowish white with a broad black band extending from behind the pronotum beneath the wings to the base of the abdomen; a large rounded black spot on the sternum just before the middle coxæ and a smaller patch just behind these coxæ; mesosternum suffused with lemonyellow. Halteres short, pale brown. Legs with the coxæ and trochanters pale testaceous; femora light yellow, the apex broadly blackened and slightly enlarged; tibiæ and tarsi dark brown, the former sometimes paler and narrowly tipped with blackish. Wings narrow, hyaline; stigmal spot brown, rounded, conspicuous.

Abdominal tergites black, the segments narrowly ringed with dull reddish yellow on the caudal margin; sternites blackish on the basal half, paler, yellowish, caudally.

Holotype, &, Dalecarlia Reservoir and outlet, District of Columbia; Aug 22, 1915 (McAtee).

Paratypes, 1♂; 1, sex uncertain, topotypic.

The type is in the collection of the United States Biological Survey, paratypes in the collection of the author.