

*Distribution*.—Massachusetts, (Harris); Ohio, (Kirtland); Lake Superior, (Agassiz); Eastern Townships, (Gosse); South side of the Gulf, (R. Bell); Rouge District.

Family IV. LYCENIDÆ Leach.

Genus 1, *Thecla*, Fab.

1. *T. Nippon*? Hübner.—Sorel and perhaps Montreal, May: (several other species undetermined.)

Genus 2, *Chrysophanus*, Hübner; *Lycæna*, Fab.; *Polyommatus*, Boisd.;

1. *C. Americana*, Harris, (American Copper).—Common, August.

*Distribution*.—Massachusetts, (Harris); Ohio, (Kirtland); Eastern Townships, (Gosse); Rouge District and Sorel.

2. *C. Thoe*? Boisd.—Lachine, (Barnston).

*Distribution*.—Rockport, Ohio, (Kirtland).

Genus 3, *Polyommatus*, Latr.; *Lycæna*, Boisd.

1. *P. pseudargiolus*, Boisd.; *Lucia*, Kirby, (Spring Azure).—Not very common, May and June.

*Distribution*.—Massachusetts, (Harris); Rockport, Ohio, (Kirtland); Eastern Townships, (Gosse); Latitude, 54° North, (Kirby); South side of the Gulf, (R. Bell); Rouge District.

2. *P. Comyntas*, Godt.—Lachine, (Barnston).

*Distribution*.—Rockport, Ohio, (Kirtland).

Family V. HESPERIDÆ, Leach.

Genus 1, *Eudamus*, Swains.

1. *E. Tityrus*, Fab.; *Clarus*, Cramer.—Numerous, 1858, (Mr. Fowler).

*Distribution*.—Massachusetts, (Harris); Ohio, (Kirtland).

Genus 2, *Pamphila*, Fab.

1. *P. cernes*, Boisd., (Tawny-edged Skipper).—Not uncommon.

*Distribution*.—Eastern Townships, (Gosse); and several undetermined species.

HETEROCERA, Boisd.

Group 1. SPHINGINA, Stainton.

Family I. SPHINGIDÆ, Leach.

Genus 1, *Smerinthus*, Latr.

1. *S. geminatus*, Say; *cerisii*, Kirby, (Twin-eyed Hawkmoth).—Not uncommon.

*Distribution*.—Massachusetts, (Harris); Orillia, C.W., (Bush); Eastern Townships (Gosse); Sorel.

2. *S. myops*, Abbot & Smith.—Rare.

*Distribution*.—United States, (Doubleday).

Genus 2, *Sphinx*, Linn.

1. *S. drupiferarum*, Abbot and Smith.—Common, June and July.

*Distribution*.—United States.