

still caused a fall of pressure, but without slowing of the heart. The fall of pressure must be in some degree independent of vagus irritation, which, however, usually accompanies it.

(15) It only remains to be considered whether the slow action or temporary stoppage of the heart with great fall of pressure produced by vagus irritation is in itself an element of danger in chloroform administration, and if it is not, wherein the danger actually lies.

(16) The experiments in which deliberate irritation of the vagi was carried on during anaesthesia show unmistakably that irritation of these nerves diminishes rather than enhances the danger of anaesthetics. The effect upon the heart is never continuous, and as the vagus becomes exhausted, or when the irritation is taken off, the blood pressure rises again, as it does when the same result is produced by asphyxia. The slowing of the heart and circulation which is produced by irritation of the vagus by any cause, such as holding the breath in chloroform administration, retards the absorption and conveyance of chloroform to the nerve centres, just as holding the breath, whether voluntary or involuntary, prevents chloroform from entering the lungs; and of itself slowing or temporary stoppage of the heart in chloroform administration is not dangerous.

(17) To answer the second part of the last question in Paragraph 15 is easy enough, if it is kept in mind that the effect of vagus irritation upon the heart is never continuous; and in chloroform administration, as the pressure rises again after the slowing of the heart and temporary fall of pressure produced by any form of asphyxia, violent respiratory efforts with bounding heart's action lead, as in the case of struggling, to a rapid and dangerous inhalation of chloroform, and consequent rapid and dangerous decline in blood pressure. It is, in fact, the temporary exhaustion of the vagi after stimulation that is to be feared, and not the actual stimulation as long as it is continued.

(18) In accordance with this fact, it will be found that in chloroform administration neither holding the breath, even if involuntary, or vagus inhibition can be kept up beyond a certain time; and if the chloroform is not removed from the face, one or both of two things may happen: (a) when the animal breathes again, it takes deep and gasping inspirations, the lungs become filled with chloroform, and an over-dose is taken in with extreme rapidity; or (b) when the restraining influence of the vagus is taken off the heart, through the irritation ceasing or the nerve becoming exhausted, the heart bounds on again, and the circulation is accelerated in proportion. The blood then becomes quickly saturated with chloroform, and an over-dose is at once conveyed to the nerve centres. The theory which has hitherto been accepted is that the danger in chloroform adminis-

tration consists in the slowing or stoppage of the heart by vagus inhibition. This is now shown to be absolutely incorrect. There is no doubt whatever that the controlling influence of the vagus on the heart is a safeguard, and that it is the exhaustion of the nerve which is dangerous.

(19) It can be readily understood how a condition in which the pulse is rapid and bounding, with high blood pressure, leads to more rapid absorption of chloroform from the lungs, and a more rapid propulsion of the chloroformed blood to the medulla oblongata, and consequently to a more rapid paralysis of the respiratory and vaso-motor centres and precipitous fall in the blood pressure. Such a condition is produced in some cases by ether or by division of both vagi or by a full dose of atropine. Not only is the poisoned blood carried more swiftly to the vital centres in these cases, but added to this there is the fact that, as the heart is already doing its utmost before the chloroform is given, it is unable to stave off by increased work the fall in pressure that occurs when the vaso-motor centre is paralysed. On the other hand, it seems clear from Experiment 92 that the direct action of chloroform upon the heart's substance is not the cause of the fall of pressure that occurs when it is inhaled.

(20) In Experiment 92 repeated injections of 20 minims of chloroform were made into the jugular vein, and its effect was not to paralyse the heart, but to produce anaesthesia and a gradual fall of blood pressure exactly as if the chloroform had been inhaled. In Experiment 72, after a considerable amount of ether had been injected into the jugular vein, and a bounding condition of pulse had been produced, the effect of injecting chloroform into the jugulars was much greater, and the fall of blood pressure much more rapid and dangerous, than in the case when chloroform alone was injected. Granting, then, the truth of Ringer's conclusions from experiments on the frog's heart (which have now been repeated and confirmed by the Commission) that chloroform has a gradual paralysing effect upon the heart's tissue, we must conclude that such an effect, in the degree in which alone it could occur in the practical inhalation of chloroform, would rather be a source of safety than of danger.

(21) The Committee discussed the advisability of cutting the vagi some time previously to experimenting on the blood pressure with chloroform. The effect of this procedure is to cause continuous rapid action and tendency to exhaustion of the heart, as well as to degeneration of the terminal branches of the nerves in the heart if the animal live sufficiently long. Such experiments might be of some interest theoretically, and also have had a practical bearing upon the condition of the heart in certain cases of chronic alcoholism; but the Committee decided not to perform them, as it con-