immediate scientific and thorough organized campaign against Venereal Disease was proven. At this Congress the widespread extent of syphilis and gonorrhea was revealed in all its horrifying magnitude: note the following statistics:

All the Canadian hospitals state that from 20% to 30% of their inmates have syphilis.

The percentage of syphilis among the recruits was 16.77.

(Vedder).

It has been estimated that 8% of the total Canadian population, 650,000 are suffering from some form of venereal disease.

(Dr. G. Bates)

The preceeding figures, large as they are, are relative to syphilis alone.

As regards gonorrhea, no reliable statistics have been computed: but, all venereal specialists are unanimous in stating that gonorrhea is from four to five times as prevalent as syphilis, and it is the most contagious disease existing, except measles.

Can a better established proof regarding the vast extent of these scourges be found?

You are aware, Gentlemen, that syphilis is not only widespread in our population, but, also it ranks first in our list of fatal diseases: it is the most used scythe in the hands of the grim Reaper. Of course, the word syphilis seldom appears on a death certificate. A milder title is employed in order to conceal the misery behind the disease. This does not alter the fact that syphilis is the cause of death in:

100% of locomotor ataxia cases.

100% of paresis cases (softening of the brain).

50% of all cases of organic troubles.

50% " angina pectoris.

40% " arterial degeneration, aneurism, etc.

40% " cerebral hemorrhage.

40% " general paralysis.

20% " " Bright's disease.

10% " epilepsy.

10% " encephalitis.

10% " meningitis.

American Social Hygiene Association. — Standard Statistics.