

- (a) VIREO OLIVACEUS—*Red-eyed Vireo*—Common.
 (a) " GILVUS—*Warbling Vireo*—Rather common.
 (a) " FLAVIFRONS—*Yellow-throated Vireo*—Rare.
 (a) MINIOTILTA VARIA—*Black and White Warbler*—Common in deep woods, but only one nest found, by W. E. Saunders in May, 1881.
 (a) HELMINTHOPHILA CHRYSOPTERA—*Golden-winged Warbler*—Rather rare.
 (a) " RUFICAPILLA—*Nashville Warbler*—Limited to a few localities; a single nest found by Wm. Saunders near London in 1882.
 (a) DENDROICA ÆSTIVA—*Yellow Warbler*—Common.
 (a) " PENNSYLVANICUS—*Chestnut-sided Warbler*—Tolerably common.
 (a) SEIURUS AUROCAPILLUS—*Ovenbird*—Tolerably common.
 (a) " NOVEBORACENSIS—*Water Thrush*—Rather rare.
 (a) GEOTHLYPIS TRICHAS—*Maryland Yellow Throat*—Common in suitable localities.
 (a) SYLVANIA CANADENSIS—*Canadian Warbler*—Rather common. Partially fledged young found in several localities near London.
 (a) SETOPHAGA RUTICILLA—*Redstart*—Common.
 (a) GALEOSCOPTES CAROLINENSIS—*Catbird*—Common.
 (a) HARPORHYNCHUS RUFUS—*Brown Thrasher*—Locally common.
 (a) TROGLODYTES AEDON—*House Wren*—Common.
 (a) TROGLODYTES HIEMALIS—*Winter Wren*—A single nest found with one egg; not uncommon in deep cedar-swamps.
 (a) CERTHIA FAMILIARIS AMERICANA—*Brown Creeper*—Rare. A nest of young found nine miles from London in June, 1880.
 (a) SITTA CAROLINENSIS—*White-breasted Nuthatch*—Not very common.
 (a) PARUS ATRICAPILLUS—*Black-capped Chickadee*—Not common.
 (a) POLIOPTILA CERULEA—*Blue-gray Gnatcatcher*—Tolerably common.
 (a) TURDUS MUSTELINUS—*Wood Thrush*—Tolerably common.
 (a) " FUSCESCENS—*Wilson's Thrush*—Common.
 (a) MERULA MIGRATORIA—*Robin*—Abundant.
 (a) SIALIA SIALIS—*Blue Bird*—Common. A few sets of white eggs have been found.

SUPPLEMENTARY LIST—No. 1.

Birds which have probably bred in greater or less numbers in the past, but have now become so rare that it is unlikely that their nests will be found:

- (b) LOPHODYTES CUCULLATUS—*Hooded Merganser*.
 (b) ECTOPISTES MIGRATORIUS—*Passenger Pigeon*.

SUPPLEMENTARY LIST—No. 2.

Birds which are observed in summer and are likely to be found breeding as the fauna is better worked up:

- (a) RALLUS VIRGINIANUS—*Virginia Rail*.
 (a) PORZANA CAROLINA—*Carolina Rail*.
 (a) GALLINULA GALEATA—*Florida Gallinule*.
 (a) SYRNIUM NEBULOSUM—*Barred Owl*—Scarce resident.
- } Will probably be found in some of our larger marshes.

(a) NYCTAL
Elgin County, in

(a) EMPID
and June.

(a) CONTO
Common summ

(a) ICTERU
observed every

(a) ZONOTE
in summer.

(a) DENDR

(a) "

(a) "

(a) "

(a) "

(a) GEOTHI

(a) CISTOTI
some of our larg

(a) REGUL
in the spruce sw

(a) TURDUS
cedar swamps.

PAL

In 1874, M
Genus Pamphil
the following s
Sassacus, Ottoe,

None of the
the differences b
sons instituted
males and two fe
should say, from
Colorado, and t
underside of sec
tion in the prom

Though no
species had been

Since then
collectors at Cac
in 1888.

The form fo
outer third of th
wings, with the
though occasiona

In 1890, on
for a day at Reg