

Beneath fore wing fuscous, with slight reddish tinge; a distinct, though not prominent, mesial band not reaching inner margin; discal dot present. Hind wings whitish outwardly, reddish-fuscous along costa and outwardly, discal dot and mesial band as on fore wing.

Type, 1 ♂. Huachuca Mts., Arizona.

*Tricholita baranca*, n. sp.—♂. Expanse, 28 mm.

General colour seal-brown, with a slight reddish tinge, more or less frosted with pale scales. Head and thorax somewhat paler, showing less of the reddish tint, owing to a greater admixture of the pale hairs. Ordinary lines all tracable in perfect specimens, though not at all prominent. Under the lens the veins seem to be somewhat darkened and very sparsely coated with pale scales. Basal half-line evident, more noticeable from the pale filling than from the only very slightly darker limiting lines. T. a. almost transverse, irregularly dentate, slightly darker than ground colour, accompanied by slightly paler inner shade. Median shade rather more prominent than the other lines, outwardly oblique to lower end of reniform, thence inwardly oblique to middle of inner margin. T. p. scalloped, only slightly exerted, beyond cell, thence quite direct to inner margin. S. t. wavy, irregular, somewhat darker than ground colour. Median and terminal spaces slightly darker than remainder of wing. Costal and basal areas somewhat more frosted than remainder of wing. The inception of the transverse lines on costa are somewhat darkened, the pale filling showing as light dots. Towards apex there are three or four more pale bands, none of which, however, are strongly pronounced, but plainly visible under the lens. Fringe concolorous with terminal space, with faint darker basal line and pale dots at end of veins. Claviform obsolete and orbicular usually so, though in one specimen a faint minute brownish ring can be made out under the lens. Reniform margined by white points, usually four in the outer row and two or three in the inner, the outer row is much better developed and the second from the costa is evidently composed of the fusion of two others, as it is about twice as large and is in some specimens partly divided. Between the two rows of pale dots the spot is filled with reddish scales. The lower point of the inner row is the largest in all the specimens before me, those lying above it showing more or less of a tendency to become obsolete. Hind wings fuscous with more or less of a reddish mixture. Very faint traces of discal dot and mesial band, scarcely discernible except in certain lights. Fringe fuscous at base, whitish externally. Beneath fore wings paler than above, darkened centrally, paler along costa and inner margin. Traces of dark extra