£1	11	3
	12	3 6
No.	7	6
1	1	8

on the first of Janubalanced, this system igation. In settling te to balance his acr give him your note ot balanced by cash

perfluous fruit that and dairy, to which third and perhaps nd cooking or sourcarried so far as to and water with the tation as adhere to nperature between Roots for fattening not intended to be small quantity of swine; then parsrtzel; ruta-bagas, e nutritive properge a bulk, that we antage. There is t cooking. When ch, to prevent imys before them .in motion, and it the most available uscle. Scraps, or r pressing the fat. ood, although not sh them off, many n slops or indifferniently procured;

steful practice.

n winds or storms,
attacks. The hog
; nature has pror of fat which surnon-conductors of
ntense cold which
ned animal.

1848. Febru'y. May	15 1	By	Cash,	:	vaca¥ oskigi	Phu Cate water	feros Isaki zhiu	8 8T	\mathcal{L}_1	2 5	6 0
6 (e f 16 (s) - 6 (kr)6										ille eletar	

paid, or note given, it should be done by giving the person credit, or charging him with enough to balance it. For instance, if you find that he is owing you, give him credit for the amount "By Balance," and charge the same to him in the new account; if, on the other hand, you find that you are owing him, make him Dr. for the amount "To Balance," and credit him with the same in the new account.

For Coughs, and Inflammation of the Lungs, bleeding should be resorted to, after which give gentle purges of Castor-oil, or Epsom Salts, which should be followed with a dose of antimonial powders—2 grains, mixed with half a drachm of nitre.

For Costiveness or Loss of Appetite, sulphur is an excellent remedy, given in a light mess.

The Itch may be cured by anointing with equal parts of lard and brimstone. Rubbing posts, and a running stream to wallow in are preventives.

The Kidney-worm is frequently fatal; and always produces weakness of the loins and hind legs, usually followed by entire prostration. Preventives, are general thrift, a range in a good pasture, and a dose of half a pint of wood-ashes every week or fortnight, in their food. A small quantity of saltpetre, spirits of turpentine or tar, will effect the same object. When attacked, apply spirits of turpentine to the loins, and administer calomel carefully, or give half a table-spoonful of copperas daily, for one or two weeks.

Blind Staggers, generally confined to pigs, shows itself by foaming at the mouth, rearing on their hind legs, champing and grinding their teeth, and apparent blindness. The best remedies are bleeding and purging freely. It is important to keep the issues on the inside of the fore-legs, just below the knee, thoroughly cleansed.

The tails of young pigs frequently drop or rot off, which may be prevented by adding a little brimstone or sulphur to the food of the dam; or rub oil or grease daily on the affected parts. It may be detected by a roughness or scabbiness at the point where separation is likely to occur.

Bleeding—the most convenient mode, is from an artery just above the knee, on the inside of the fore-arm; it may be drawn more copiously from the roof of the mouth. Hogs confined in styes, should have a few shovels-ful of soil, or rotten wood, thrown to them weekly, and be supplied with salt daily. Plenty of clean, dry straw, and nourishing food, are indispensable in the winter, and will abundantly repay all cost.

The Hedgehog —It has been ascertained that the most violent poisons have no effect on the hedgehog; and consequently the animal must be very useful in the forests, etc. for destroying noxious reptiles.