1908, 2 ?'s. Summit of Sulphur Mt., Aug. 21, 31, 1908, 2 ?'s; Aug, 23, 1909, 1 ?. Mt. Inglesmaldie, 6,000 to 7,000 ft., Sept. 11, 1908, 1 ?, 3 ?'s. (Sanson.) Also recorded from Laggan (Scudder⁸), Banff (Walker), and Edmonton (Fletcher²). The only record of the macropterous form is the one from Edmonton. This form is common in northern Ontario.

80. Melanoplus femur-rubrum (De Geer).

MAN.—Winnipeg. (Scudder⁵.) Portage la Prairie and Brandon. (Walker.)

SASK.—Condee, 1 &; Findlater, 2 & 's; Pasqua, 1 &; Weyburn, 2 & 's. (Alexander.) Also recorded from Moose Jaw and Waldeck (Walker).

ALTA.—Ponoka, 1 9. (Alexander.) Walsh, Aug. 23, 1901,3 & s, 2 9 's. (Willing.) Also recorded from Medicine Hat (Caudell³).

B. C.—(Bruner¹, Scudder⁵.) Vancouver Id. (Scudder⁵.) Agassiz. (Walker.) Kaslo. (Caudell⁴.)

The specimens from the drier parts of Saskatchewan are considerably below the medium size.

81. Melanoplus extremus (Walker).

MAN.—Aweme, June 20, 1905, 1 &. (Criddle.) Also recorded from Plum Coulee (Walker) and Aweme (Fletcher³).

SASK.—Kinistino, June 23, 1905, 1 &, 1 \(\frac{7}{3}, \) (teneral). (Willing.) ALTA.—Banff, July 22, 31, 1908, 2 & is; Aug. 5, 28, 29, 1908, 4 & is, 3 \(\frac{7}{3} \) s. (Sanson.) Also recorded from Beaver Lake (Fletcher⁸) and Banff (Scudder⁵).

B. C.-Field. (Rehn.)

The only macropterous individuals in the series are two males from Banff.

82. Melanoplus monticola Bruner.

ALTA.—Mountains near Laggan. (Caudell3.)

83. Melanoplus angustipennis (Dodge).

MAN.—Aweme, July 19, 1904, 1 &; July 29, 1906, 1 &; Aug. 12, 1905, 1 &; Aug. 18, 1906, 1 &; Aug. 23, 1904, 1 &; Oct. 6, 1907, 1 &. (Criddle.)

The specimen taken Aug. 18, 1906, has blue hind tibiæ, the others red. The red-legged individuals, commonly known as *M. coccineipes*, are certainly not distinct from the typical bluelegged form.