ections, 472, , actiology, isillitis (cone tissue of esthesia in, e of hyoid and cricoid ge, 557 586 lue to scald, on septal igerous, 27 30 onset of, 31 475 used in, 25 of, in exparesis in, rhinitis, 31 481 alysis in. 545 thyro arypharynx, of smell bil..teral iti- 565

f Jarynx.

Interarytenoid fold, 517 folds, swelling in influenza, 482 pachydermia, 560 region, pachydermia of, 571 Interarytenoideus affected by tuberculosis of larynx, 727 paralysis of, 727 Intranasal disease, ocular affections accompanying, cause, 68 operation, caution against wounding mucous membrane surfaces in close apposition, 42 route in treatment of empyema of antrum, 220 treatment failing to relieve symptoms of reflex origin, 68 Intrathoracic tumour cause of unilateral abductor paralysis, 722 Intratonsillar or tonsillar fossa, 396, 397, 398 Intratracheal injections in relief of cough in laryngeal tuberculosis, 643 Intubation and tracheotomy, comparative mcrits of, 674, 677 relative merits of, in chronic stenosis of larynx, 678 for chronic laryngeal stenosis, necessity of supervision of patient after removal of tube, 679 in acute stenosis of larynx, 670 in adults, difficulties of, 679 in cancer of æsophagus, 754 in chronic stenosis of larynx, 678 in diphtheria, 495 in inherited syphilis of larynx, 661 in laryngeal diphtheria, advantages of, 675 difficulty of feeding in, 676 disadvantages, 675 preference of, to tracheotomy for children under three years, 675 papillomata in children, 604 stenosis, 461 death caused by removal of tube, case, 679 in membranous laryngitis, 554 in spasmodic laryngitis, 548 instruments for. O'Dwyer's, 670, length of time tube required in, 677 method of performance, 671, 672 Iodine, nascent, Pfannenstiel's method of treatment by, in nasal tuberculosis, 150 solution, application in pachydermia laryngis, 573 lonisation in atrophic rhinitis, 54 lion and strychnine in treatment of diphtheritic laryngeal anæsthesia, 731

815 Iron and strychnine, in acute laryngitis, 545 in tonsillitis (convalescent stage), in paralysis of palato-pharyngeal muscles of toxic origin, 393 in sensory neuroses of pharynx, 395 in syphilis, 164 iodide of, syrup of, in laryngismus stridulus, 707 perchloride of, in acute tonsillitis, 411 application in chronic laryngitis, 563 to larynx in functional aphonia, 719 preparations of, in functional aphonia (bilateral adductor paralysis), 718 sulphate of, and sulphate of magnesium in prevention of recurrence of epistaxis, 184 valerianate, zinc, and quinine, in swelling of erectile tissue of nose, 70 Irrigation, dangers of, in exploration of antrum, 215 nasal, 295 Irrigations in acute suppurative inflammation of antrum, 204 Jackson, Chevalie:, avoidance of anæsthesia in peroral endoscopy, 524 Jaenicke polypus snare, 128 Jansen's bone forceps, 274 Karyorrhexis, 6 Keith, A., F.R.S., traction diverticula of cesophagus, 767 Kelly, A. Brown, cystic disease of antrum, 225 difficulties and dangers of exploratory puncture of antrum of Highmorc, 215, 216 naso-antral or choanal polypus, 138 rare form of congenital syphilis of plarynx and larynx, 384

Kernig's sign in cerebro-spinal fever, 502 Kidd, Percy, form of laryngeal tuberculosis simulating bilateral abductor paralysis, 591 Kidney, granular contracted, and hypertrophy of heart, occurrence of epistaxis in, 182 Killian's apparatus for suspension laryngoscopy, 527, 528 long-bladed nasal speculum, 16, 107 operation, 276 Kirstein's forehead lamp, 521