

1912. *R. MARITIMUS*, L. (Golden Dock.)

River bank. Rare. Green's Creek (*A. H. Moore*). New Edinburgh (*H. M. Ami*). Hull (*Prof. Macoun*). Aug.—2. A low plant, with crowded, leafy, compact spikes of yellow fruit. Seed-valves spear-shaped, with two long teeth on each side and a large grain on the back.

1913. *R. Acetosa*, L. (Common Sorrel.)

Occasionally introduced with grass seed; but not persistent. Billings Bridge. Hintonburgh. July—2. Dioecious. Seed-valves grain-bearing, enlarging as the fruit ripens.

1915. *R. ACETOSELLA*, L. (Sheep Sorrel.)

Abundant in sandy soil and exhausted fields. July—1. Dioecious. Seed-valves ovate, scarcely enlarging in fruit, not grain-bearing.

## ARISTOLOCHACEÆ.

## ASARUM, L. Asarabacca.

1916. *A. CANADENSE*, L. (Wild Ginger.)

Rich woods. Common. May—4. (B) A charming plant, with its rich purplish brown, three-cleft flower, borne low on the ground beneath the leaves, in the axil of the two delicate green, pubescent, kidney-shaped leaves. The fleshy root is aromatic, when broken smelling like ginger.

## PIPERACEÆ.—Pepper Family.

## SAURURUS, L. Lizard's Tail.

1918. *S. CERNUUS*, L.

In shallow water. In great abundance along the Nation River at Casselman. July—3. This is the only station so far recorded in this part of Canada.

## THYMELEACEÆ. Mezereon Family.

## DAPHNE, L.

1921. *D. Mezereum*, L. (Mezereon.)

Rocky woods. Near Hemlock Lake, Beechwood. (Mrs. Chan. rlin.) Ap.—4. Introduced, but well established. The beautiful pink or white fragrant flowers opening before the leaves. Followed by scarlet berries.