

Safety pins, one package.  
 Plain pins, one package.  
 Soap.  
 Adhesive plaster, two yards.  
 Tourniquet, rubber, one.  
 One pair scissors, ordinary.  
 One pair scissors, surgical.  
 One jack knife.  
 One razor.

Three pairs artery forceps.  
 The dimensions of this box are 15x11x7 $\frac{1}{2}$  inches. It is set inside a larger box which contains two pair blankets, a hot water bottle, and a number, say fifteen, half and quarter-inch boards of assorted lengths and widths. One or two boards about five feet long and four inches wide should be with the box to use in case of fracture of the thigh. In smaller industries it would not be wise to have such a large supply on hand as it deteriorates with age, but loss can be prevented by filling the small boxes in the factory from the large box and have it in turn replenished from the drug store.

Two pairs of crutches and two stretchers are a very necessary part of the equipment. The ambulance men should be trained and practised in stretcher drill until they are proficient.

Chairman,—

We have all heard Dr. Cannon's very able lecture and I am sure that if there are points that he has mentioned that you would like further enlightenment on, that Dr. Cannon will be only too pleased to help you out and clear up any difficulties you may have.

Mr. Beswick,—

I would like to ask the doctor what time would elapse after constricting a limb before gangrene would set in.

Dr. Cannon,—

That depends on how completely the circulation is shut off. However, it would not be for some time in any case. Fortunately it is almost impossible to completely shut off the flow of blood, for instance the bone carries a certain amount of blood. I do not think you need ever let that worry you, if you can get a doctor within an hour or two the limb would easily recover.

Mr. Baldwin,

In case of compound fracture. You remarked about the