

SAPERDA FAYI, S. CONCOLOR AND APHODIUS RUFIPES.

BY JOHN HAMILTON, M. D., ALLEGHENY, PA.

SAPERDA FAYI, Bland.—This beautiful Saperda breeds in the small limbs of Crataegus, especially *crus-galli* and *tomentosa*, as first observed by Mr. C. D. Zimmermann, CAN. ENT., 10, 220; and should it, like some of its allies, acquire a taste for cultivated fruit trees, it would be a formidable enemy, as is evidenced by the way it depredates on thorn bushes. The beetles appear here the last week in May or the first week in June, according to the season, the males preceding the females three or four days. They do not appear to eat and are short lived, the whole brood (except stragglers) appearing and disappearing within the space of ten or twelve days, so that should the collector be negligent, or the weather unsuitable for collecting at the time of their appearance, he may get none till the next season. As soon as the females appear the males are ready to associate with them, the union lasting three or four hours. They are not much given to flying about, usually ovipositing on the same tree they inhabited as larvæ. There may be several thorn trees not far apart, and one will be depredated on year after year till it is nearly destroyed, while the others will remain untouched till colonized apparently by accident. The beetles are sluggish, and when approached suddenly fall to the ground and quickly endeavor to conceal themselves, not feigning death, as many insects under the same circumstances do; and when I say feigning death, I mean it literally, in opposition to an unsupported dogmatic statement which I lately saw in print somewhere, "that insects can have no knowledge of death."

Oviposition is effected probably during the night, and the process has not been witnessed, nor the eggs seen. The limbs selected for this purpose vary from one third to one and one fourth inches in diameter, and according to the thickness of the limb, the female with her powerful mandibles makes from three to six longitudinal incisions through the bark, each about three fourths of an inch long and equi-distant and parallel to one another, dividing the circumference into sections nearly equal; an egg is placed in each end of each of these slits, and as soon as hatched the larva makes a burrow beneath the outer layer of wood, perhaps one eighth inch in length at first, and uses this as a retreat whence it issues to feed on the diseased wood caused by the incision. These slits and the