

Sp. Pter. maculipennis. Fem. *Viridis*, abdomen cupreum, antennæ nigrae, pedes flavi, femora fusca, proalæ fusco maculatæ.

"Viridis, parum nitens: oculi ocellique rufi: antennæ nigrae; articulus 1^{us} flavus; 2^{us}, 3^{us} et 4^{us} fusci: abdomen cupreum; segmentum 1^{um} laete cupreo-æneum; oviductus rufus; pedes flavi: coxæ virides; femora fusca; meso- et metatarsi pallide flavi, apice fusci: alæ sublimpidæ; prealæ sub nervis ulnari et cubitali fuscomaculatæ; squamulæ fusca; nervi fulvi; stigma minutum. (Corp. long. lin. 1-1 $\frac{1}{4}$; alar lin. 1 $\frac{1}{4}$ -1 $\frac{1}{2}$.)

"Var. B.—Abdomen purpureo-cupreum; segmentum 1^{um} laete cupreoviride: femora flava

"Var. c. Var. b. similis: caput et thorax viridi-ænea.

"July, August: grass in fields; near London."

De Dalla Torre, 1898, p. 134.

2. *Arthrolytus albiscapus* Thomson.

Thomson, 1878, pp. 159-160.

Möller, 1882, p. 180.

De Dalla Torre, 1898, p. 155.

Schmiedeknecht, 1909, p. 359.

"2. *A. albiscapus* m: Viridis pedibus flavis, coxis posticis supra cyaneis tibiis antennisque scape albidis, abdomine aureo, alis macula magna disci fusca. ♀ Long., 3-4 mill.

"Præcedenti statura similis, sed facie infra medium valde protuberante, antennis scapo longiore, albo, flagello validiore sed haud incrassato, thorace laetius viridi subtilissime punctato, collari multo augustiore, metathorace brevior et magis declivi, costula nulla, alis limpidis, radio subangulo^a acutior egrediente, stigmatibus crassioribus, pedibus præsertim femoribus adhuc validioribus, mandibulis externe sinuatis, apice forcipatis bene distinctus.

"Sällsynt i medlersta och södra Sverige."^b

(Thomson, 178, pp. 159-160.)

This species is mentioned by Möller (1882) in his description of *puncticollis*. With this exception, it is not again mentioned in the literature excepting in the citations given.

a. Beginning p. 160.

b. R rely, in middle and southern Sweden.